

For Reference

NOT TO BE TAKEN FROM THIS ROOM

For Reference

NOT TO BE TAKEN FROM THIS ROOM

Ex libris
UNIVERSITATIS
ALBERTAENSIS





Digitized by the Internet Archive
in 2019 with funding from
University of Alberta Libraries

<https://archive.org/details/Schuld1967>

THE UNIVERSITY OF ALBERTA

RAPESEED MEAL AS A PROTEIN SOURCE FOR SWINE REPRODUCTION

by

FREDERICK WILLIAM SCHULD

A THESIS

SUBMITTED TO THE FACULTY OF GRADUATE STUDIES

IN PARTIAL FULFILMENT OF THE REQUIREMENTS FOR THE DEGREE OF

MASTER OF SCIENCE

DEPARTMENT OF ANIMAL SCIENCE

EDMONTON, ALBERTA

APRIL, 1967

THE UNIVERSITY OF CHICAGO

DEPARTMENT OF THE HISTORY OF ARTS AND ARCHITECTURE

RESEARCH REPORT

1981

RESEARCH REPORT ON THE HISTORY OF ARTS AND ARCHITECTURE

RESEARCH REPORT ON THE HISTORY OF ARTS AND ARCHITECTURE

RESEARCH REPORT

RESEARCH REPORT

RESEARCH REPORT

RESEARCH REPORT

UNIVERSITY OF ALBERTA
FACULTY OF GRADUATE STUDIES

The undersigned certify that they have read, and recommend to the Faculty of Graduate Studies for acceptance, a thesis entitled "Rapeseed Meal as a Protein Source for Swine Reproduction" submitted by Frederick William Schuld, B.Sc., in partial fulfilment of the requirements for the degree of Master of Science.

ABSTRACT

Experiments were designed to study the effects on growth, reproduction and lactation and on growth of progeny when 8% solvent-extracted rapeseed meal was substituted for an isonitrogenous level of soybean meal in the diets. Two experiments were conducted: one with sows and one with their offspring. Experiment 1, using 24 sows, was factorially designed with 2 levels of dietary rapeseed meal (0 and 8%) to 90 kg liveweight with each lot further subdivided between 0 and 8% rapeseed meal for reproduction. In experiment 2, offspring of both sexes from the dams receiving either 0 or 8% rapeseed meal were allotted to either 0 or 8% rapeseed meal diets from weaning to market weight.

Feeding rapeseed meal to gilts tended to depress rate of gain in the period from 55 to 90 kg liveweight and in the pre-gestation period. Weaning weights of litters from sows fed rapeseed meal during growth were depressed. The fat content of colostrum was higher in sows fed rapeseed meal during growth. The increased fat content was reflected in a rise in total solids content of the colostrum. The fatty acid composition of the colostrum and milk fat was not affected by dietary treatments. The feeding of rapeseed meal during gestation and lactation had no significant effects on sow or litter performance.

In experiment 2, the feeding of rapeseed meal led to a depression in feed consumption and in gain in the finishing period from 55 to 90 kg liveweight and a depression in efficiency of feed utilization from weaning to market weight. No differences were observed in the digestibility and retention of energy and nitrogen of pigs fed either the basal soybean meal diets or the rapeseed meal diets. The carcasses of pigs fed rapeseed meal were leaner based on reduced backfat depth, increased ham weight as a percent of carcass weight and a trend toward higher Record of Performance scores.

THE UNIVERSITY OF CHICAGO PRESS

100 N. LAUREL STREET, CHICAGO, ILL. 60602

1980

100 N. LAUREL STREET, CHICAGO, ILL. 60602

100 N. LAUREL STREET, CHICAGO, ILL. 60602

100 N. LAUREL STREET, CHICAGO, ILL. 60602

100 N. LAUREL STREET, CHICAGO, ILL. 60602

100 N. LAUREL STREET, CHICAGO, ILL. 60602

100 N. LAUREL STREET, CHICAGO, ILL. 60602

100 N. LAUREL STREET, CHICAGO, ILL. 60602

100 N. LAUREL STREET, CHICAGO, ILL. 60602

100 N. LAUREL STREET, CHICAGO, ILL. 60602

100 N. LAUREL STREET, CHICAGO, ILL. 60602

100 N. LAUREL STREET, CHICAGO, ILL. 60602

100 N. LAUREL STREET, CHICAGO, ILL. 60602

100 N. LAUREL STREET, CHICAGO, ILL. 60602

100 N. LAUREL STREET, CHICAGO, ILL. 60602

100 N. LAUREL STREET, CHICAGO, ILL. 60602

100 N. LAUREL STREET, CHICAGO, ILL. 60602

100 N. LAUREL STREET, CHICAGO, ILL. 60602

100 N. LAUREL STREET, CHICAGO, ILL. 60602

100 N. LAUREL STREET, CHICAGO, ILL. 60602

100 N. LAUREL STREET, CHICAGO, ILL. 60602

100 N. LAUREL STREET, CHICAGO, ILL. 60602

100 N. LAUREL STREET, CHICAGO, ILL. 60602

100 N. LAUREL STREET, CHICAGO, ILL. 60602

100 N. LAUREL STREET, CHICAGO, ILL. 60602

ACKNOWLEDGMENTS

I wish to thank Dr. L.W. McElroy, Head of the Department and Professor of Animal Science, for placing the facilities of the Department at my disposal. I gratefully acknowledge the valuable advice and assistance given during this study by Dr. J.P. Bowland, Professor of Animal Nutrition, and his helpful suggestions and criticisms for the preparation and editing of this manuscript.

Thanks is expressed to Mr. G. Stephens, swine herdsman at the University of Alberta Livestock Farm, and his staff for their care of the experimental swine and their assistance and willing cooperation in this study.

I gratefully acknowledge the patience, assistance and encouragement of my wife, Jennie, in the course of this study. I thank Miss Sandra Popik for her assistance in typing this manuscript.

Financial support provided by grants from the National Research Council of Canada and the Alberta Agricultural Research Trust is acknowledged.

1. The first part of the paper is devoted to the study of the

properties of the function $f(x)$ defined by the equation

$$f(x) = \int_0^x \frac{1}{1+t^2} dt$$

and to the study of the function $g(x)$ defined by the equation

$$g(x) = \int_0^x \frac{1}{1+t^2} dt$$

and to the study of the function $h(x)$ defined by the equation

$$h(x) = \int_0^x \frac{1}{1+t^2} dt$$

and to the study of the function $i(x)$ defined by the equation

$$i(x) = \int_0^x \frac{1}{1+t^2} dt$$

and to the study of the function $j(x)$ defined by the equation

$$j(x) = \int_0^x \frac{1}{1+t^2} dt$$

and to the study of the function $k(x)$ defined by the equation

$$k(x) = \int_0^x \frac{1}{1+t^2} dt$$

and to the study of the function $l(x)$ defined by the equation

$$l(x) = \int_0^x \frac{1}{1+t^2} dt$$

and to the study of the function $m(x)$ defined by the equation

$$m(x) = \int_0^x \frac{1}{1+t^2} dt$$

and to the study of the function $n(x)$ defined by the equation

$$n(x) = \int_0^x \frac{1}{1+t^2} dt$$

and to the study of the function $o(x)$ defined by the equation

$$o(x) = \int_0^x \frac{1}{1+t^2} dt$$

and to the study of the function $p(x)$ defined by the equation

$$p(x) = \int_0^x \frac{1}{1+t^2} dt$$

and to the study of the function $q(x)$ defined by the equation

$$q(x) = \int_0^x \frac{1}{1+t^2} dt$$

and to the study of the function $r(x)$ defined by the equation

$$r(x) = \int_0^x \frac{1}{1+t^2} dt$$

and to the study of the function $s(x)$ defined by the equation

$$s(x) = \int_0^x \frac{1}{1+t^2} dt$$

TABLE OF CONTENTS

	Page
INTRODUCTION -----	1
REVIEW OF LITERATURE -----	2
A. Goitrogenic Properties of Rapeseed Meal -----	2
1. Nature of the potentially toxic factors -----	2
2. Action of goitrogens -----	3
B. Nutritive Value of Rapeseed Meal -----	5
1. Effect of processing methods -----	5
2. Value in starting diets -----	6
3. Value in growing and finishing diets -----	7
C. Reproduction and Lactation -----	9
D. Composition of Sows' Colostrum and Milk -----	10
1. Gross composition -----	10
2. Fatty acid composition of milk fat -----	12
EXPERIMENTAL -----	15
A. Objectives -----	15
B. Methods and Procedures -----	15
1. General -----	15
2. Formulation of experimental diets and methods of feeding -----	16
3. Experiment 1 -----	17
a. Sow growth and reproduction experiments ----	17
b. Lactation experiments -----	21
4. Experiment 2 -- Growth and metabolism experiments--	22
5. Methods of chemical analysis -----	24
6. Methods of statistical analysis -----	26

Table of Contents

1	Introduction	1
2	Chapter 1: The History of the Book	2
3	Chapter 2: The Structure of the Book	3
4	Chapter 3: The Language of the Book	4
5	Chapter 4: The Style of the Book	5
6	Chapter 5: The Content of the Book	6
7	Chapter 6: The Form of the Book	7
8	Chapter 7: The Function of the Book	8
9	Chapter 8: The Value of the Book	9
10	Chapter 9: The Future of the Book	10
11	Chapter 10: The Conclusion of the Book	11
12	Appendix A: The Bibliography	12
13	Appendix B: The Index	13
14	Appendix C: The Glossary	14
15	Appendix D: The List of Figures	15
16	Appendix E: The List of Tables	16
17	Appendix F: The List of References	17
18	Appendix G: The List of Abbreviations	18
19	Appendix H: The List of Symbols	19
20	Appendix I: The List of Equations	20
21	Appendix J: The List of Diagrams	21
22	Appendix K: The List of Maps	22
23	Appendix L: The List of Photographs	23
24	Appendix M: The List of Audio Clips	24
25	Appendix N: The List of Video Clips	25
26	Appendix O: The List of Interactive Elements	26
27	Appendix P: The List of Supplementary Materials	27
28	Appendix Q: The List of Endnotes	28
29	Appendix R: The List of Footnotes	29
30	Appendix S: The List of References	30
31	Appendix T: The List of Bibliography	31
32	Appendix U: The List of Index	32
33	Appendix V: The List of Glossary	33
34	Appendix W: The List of Figures	34
35	Appendix X: The List of Tables	35
36	Appendix Y: The List of References	36
37	Appendix Z: The List of Abbreviations	37
38	Appendix AA: The List of Symbols	38
39	Appendix AB: The List of Equations	39
40	Appendix AC: The List of Diagrams	40
41	Appendix AD: The List of Maps	41
42	Appendix AE: The List of Photographs	42
43	Appendix AF: The List of Audio Clips	43
44	Appendix AG: The List of Video Clips	44
45	Appendix AH: The List of Interactive Elements	45
46	Appendix AI: The List of Supplementary Materials	46
47	Appendix AJ: The List of Endnotes	47
48	Appendix AK: The List of Footnotes	48
49	Appendix AL: The List of References	49
50	Appendix AM: The List of Bibliography	50
51	Appendix AN: The List of Index	51
52	Appendix AO: The List of Glossary	52
53	Appendix AP: The List of Figures	53
54	Appendix AQ: The List of Tables	54
55	Appendix AR: The List of References	55
56	Appendix AS: The List of Abbreviations	56
57	Appendix AT: The List of Symbols	57
58	Appendix AU: The List of Equations	58
59	Appendix AV: The List of Diagrams	59
60	Appendix AW: The List of Maps	60
61	Appendix AX: The List of Photographs	61
62	Appendix AY: The List of Audio Clips	62
63	Appendix AZ: The List of Video Clips	63
64	Appendix BA: The List of Interactive Elements	64
65	Appendix BB: The List of Supplementary Materials	65
66	Appendix BC: The List of Endnotes	66
67	Appendix BD: The List of Footnotes	67
68	Appendix BE: The List of References	68
69	Appendix BF: The List of Bibliography	69
70	Appendix BG: The List of Index	70
71	Appendix BH: The List of Glossary	71
72	Appendix BI: The List of Figures	72
73	Appendix BJ: The List of Tables	73
74	Appendix BK: The List of References	74
75	Appendix BL: The List of Abbreviations	75
76	Appendix BM: The List of Symbols	76
77	Appendix BN: The List of Equations	77
78	Appendix BO: The List of Diagrams	78
79	Appendix BP: The List of Maps	79
80	Appendix BQ: The List of Photographs	80
81	Appendix BR: The List of Audio Clips	81
82	Appendix BS: The List of Video Clips	82
83	Appendix BT: The List of Interactive Elements	83
84	Appendix BU: The List of Supplementary Materials	84
85	Appendix BV: The List of Endnotes	85
86	Appendix BW: The List of Footnotes	86
87	Appendix BX: The List of References	87
88	Appendix BY: The List of Bibliography	88
89	Appendix BZ: The List of Index	89
90	Appendix CA: The List of Glossary	90
91	Appendix CB: The List of Figures	91
92	Appendix CC: The List of Tables	92
93	Appendix CD: The List of References	93
94	Appendix CE: The List of Abbreviations	94
95	Appendix CF: The List of Symbols	95
96	Appendix CG: The List of Equations	96
97	Appendix CH: The List of Diagrams	97
98	Appendix CI: The List of Maps	98
99	Appendix CJ: The List of Photographs	99
100	Appendix CK: The List of Audio Clips	100

	Page
RESULTS AND DISCUSSION -----	27
A. Growth, Reproduction and Lactation of Sows --	
Experiment 1 -----	27
1. Feed consumption, daily gain, and efficiency of feed utilization during pre-gestation -----	27
a. Average daily feed -----	27
b. Average daily gain -----	29
c. Efficiency of feed utilization -----	30
2. Reproductive performance -----	31
a. First litter -----	31
b. Second litter -----	35
c. First and second litters -----	36
3. Lactation studies -----	36
a. Gross colostrum and milk composition -----	41
b. Fatty acid composition -----	41
B. Progeny Growth Studies -- Experiment 2 -----	42
1. Feed consumption, daily gain and efficiency of feed utilization -----	42
a. Average daily feed consumption -----	42
b. Average daily gain -----	43
c. Efficiency of feed utilization -----	46
2. Carcass studies -----	47
C. Digestibility and Retention of Energy and Nitrogen -	50
1. Starting diets -----	50
2. Growing diets -----	51
GENERAL SUMMARY AND CONCLUSIONS -----	52
BIBLIOGRAPHY -----	56
APPENDIX	

PUBLISHED WEEKLY
535 N. Dearborn Ave., Chicago, Ill., U.S.A.

Subscription price, Five Dollars Per Annum in Advance
Single Copies, Fifteen Cents

Entered as Second-Class Matter, June 26, 1902
Postpaid

Acceptance for mailing at special rate of postage provided for in
Act of October 3, 1917. Authorized by Act of October 3, 1917.

Postmaster: Please send address changes in advance of publication.

Copyright, 1938, by American Medical Association

Printed at the American Medical Association, 535 N. Dearborn Ave., Chicago, Ill.

Second-class postage paid at Chicago, Ill., and at additional mailing offices.

Postage paid at Chicago, Ill., and at additional mailing offices.

Postage paid at Chicago, Ill., and at additional mailing offices.

Postage paid at Chicago, Ill., and at additional mailing offices.

Postage paid at Chicago, Ill., and at additional mailing offices.

Postage paid at Chicago, Ill., and at additional mailing offices.

Postage paid at Chicago, Ill., and at additional mailing offices.

Postage paid at Chicago, Ill., and at additional mailing offices.

Postage paid at Chicago, Ill., and at additional mailing offices.

Postage paid at Chicago, Ill., and at additional mailing offices.

Postage paid at Chicago, Ill., and at additional mailing offices.

Postage paid at Chicago, Ill., and at additional mailing offices.

Postage paid at Chicago, Ill., and at additional mailing offices.

Postage paid at Chicago, Ill., and at additional mailing offices.

Postage paid at Chicago, Ill., and at additional mailing offices.

Postage paid at Chicago, Ill., and at additional mailing offices.

Postage paid at Chicago, Ill., and at additional mailing offices.

Postage paid at Chicago, Ill., and at additional mailing offices.

LIST OF TABLES

	Page
Table 1. Formulation of the diets on an air-dry basis -----	18
Table 2. Composition of the diets on an air-dry basis -----	19
Table 2A. Composition of rapeseed meal compared to soybean meal -	20
Table 3. Allotment of gilts to experimental diets-experiment 1 -	20
Table 4. Breeding schedule -----	21
Table 5. Allotment of progeny to experimental diets-experiment 2	24
Table 6. Average daily feed, average daily gain, and efficiency of feed utilization -- experiment 1 -----	28
Table 7. Treatment means for average daily feed, average daily gain and efficiency of feed utilization -- experiment 1	
Table 8. Reproductive performance -- first litters, -- experiment 1 --- -----	32
Table 9. Reproductive performance -- second litters, -- experiment 1 -----	33
Table 10. Treatment means for reproductive performance -----	34
Table 11. Gross composition of colostrum and milk -----	37
Table 12. Treatment means for gross composition of colostrum and milk -----	38
Table 13. Fatty acid composition of sows' colostral and milk fat -----	39
Table 14. Treatment means for fatty acid composition of colostral and milk fat -----	40
Table 15. Average daily feed, average daily gain and efficiency of feed utilization -- experiment 2 -----	44
Table 16. Treatment means for average daily feed, average daily gain and efficiency of feed utilization -- experiment 2 -----	45
Table 17. Carcass measurements -- experiment 2 -----	48
Table 18. Treatment means for carcass measurements -----	48
Table 19. Digestibility and retention studies -----	49
Table 20. Treatment means for digestibility studies -----	49

TABLE OF CONTENTS

1	Introduction	1
2	Chapter I	2
3	Chapter II	3
4	Chapter III	4
5	Chapter IV	5
6	Chapter V	6
7	Chapter VI	7
8	Chapter VII	8
9	Chapter VIII	9
10	Chapter IX	10
11	Chapter X	11
12	Chapter XI	12
13	Chapter XII	13
14	Chapter XIII	14
15	Chapter XIV	15
16	Chapter XV	16
17	Chapter XVI	17
18	Chapter XVII	18
19	Chapter XVIII	19
20	Chapter XIX	20
21	Chapter XX	21
22	Chapter XXI	22
23	Chapter XXII	23
24	Chapter XXIII	24
25	Chapter XXIV	25
26	Chapter XXV	26
27	Chapter XXVI	27
28	Chapter XXVII	28
29	Chapter XXVIII	29
30	Chapter XXIX	30
31	Chapter XXX	31
32	Chapter XXXI	32
33	Chapter XXXII	33
34	Chapter XXXIII	34
35	Chapter XXXIV	35
36	Chapter XXXV	36
37	Chapter XXXVI	37
38	Chapter XXXVII	38
39	Chapter XXXVIII	39
40	Chapter XXXIX	40
41	Chapter XL	41
42	Chapter XLI	42
43	Chapter XLII	43
44	Chapter XLIII	44
45	Chapter XLIV	45
46	Chapter XLV	46
47	Chapter XLVI	47
48	Chapter XLVII	48
49	Chapter XLVIII	49
50	Chapter XLIX	50
51	Chapter L	51
52	Chapter LI	52
53	Chapter LII	53
54	Chapter LIII	54
55	Chapter LIV	55
56	Chapter LV	56
57	Chapter LVI	57
58	Chapter LVII	58
59	Chapter LVIII	59
60	Chapter LIX	60
61	Chapter LX	61
62	Chapter LXI	62
63	Chapter LXII	63
64	Chapter LXIII	64
65	Chapter LXIV	65
66	Chapter LXV	66
67	Chapter LXVI	67
68	Chapter LXVII	68
69	Chapter LXVIII	69
70	Chapter LXIX	70
71	Chapter LXX	71
72	Chapter LXXI	72
73	Chapter LXXII	73
74	Chapter LXXIII	74
75	Chapter LXXIV	75
76	Chapter LXXV	76
77	Chapter LXXVI	77
78	Chapter LXXVII	78
79	Chapter LXXVIII	79
80	Chapter LXXIX	80
81	Chapter LXXX	81
82	Chapter LXXXI	82
83	Chapter LXXXII	83
84	Chapter LXXXIII	84
85	Chapter LXXXIV	85
86	Chapter LXXXV	86
87	Chapter LXXXVI	87
88	Chapter LXXXVII	88
89	Chapter LXXXVIII	89
90	Chapter LXXXIX	90
91	Chapter LXXXX	91
92	Chapter LXXXXI	92
93	Chapter LXXXXII	93
94	Chapter LXXXXIII	94
95	Chapter LXXXXIV	95
96	Chapter LXXXXV	96
97	Chapter LXXXXVI	97
98	Chapter LXXXXVII	98
99	Chapter LXXXXVIII	99
100	Chapter LXXXXIX	100
101	Chapter LXXXXX	101
102	Chapter LXXXXXI	102
103	Chapter LXXXXXII	103
104	Chapter LXXXXXIII	104
105	Chapter LXXXXXIV	105
106	Chapter LXXXXXV	106
107	Chapter LXXXXXVI	107
108	Chapter LXXXXXVII	108
109	Chapter LXXXXXVIII	109
110	Chapter LXXXXXIX	110
111	Chapter LXXXXXX	111
112	Chapter LXXXXXXI	112
113	Chapter LXXXXXXII	113
114	Chapter LXXXXXXIII	114
115	Chapter LXXXXXXIV	115
116	Chapter LXXXXXXV	116
117	Chapter LXXXXXXVI	117
118	Chapter LXXXXXXVII	118
119	Chapter LXXXXXXVIII	119
120	Chapter LXXXXXXIX	120
121	Chapter LXXXXXXX	121
122	Chapter LXXXXXXXI	122
123	Chapter LXXXXXXXII	123
124	Chapter LXXXXXXXIII	124
125	Chapter LXXXXXXXIV	125
126	Chapter LXXXXXXXV	126
127	Chapter LXXXXXXXVI	127
128	Chapter LXXXXXXXVII	128
129	Chapter LXXXXXXXVIII	129
130	Chapter LXXXXXXXIX	130
131	Chapter LXXXXXXXI	131
132	Chapter LXXXXXXXII	132
133	Chapter LXXXXXXXIII	133
134	Chapter LXXXXXXXIV	134
135	Chapter LXXXXXXXV	135
136	Chapter LXXXXXXXVI	136
137	Chapter LXXXXXXXVII	137
138	Chapter LXXXXXXXVIII	138
139	Chapter LXXXXXXXIX	139
140	Chapter LXXXXXXXI	140
141	Chapter LXXXXXXXII	141
142	Chapter LXXXXXXXIII	142
143	Chapter LXXXXXXXIV	143
144	Chapter LXXXXXXXV	144
145	Chapter LXXXXXXXVI	145
146	Chapter LXXXXXXXVII	146
147	Chapter LXXXXXXXVIII	147
148	Chapter LXXXXXXXIX	148
149	Chapter LXXXXXXXI	149
150	Chapter LXXXXXXXII	150
151	Chapter LXXXXXXXIII	151
152	Chapter LXXXXXXXIV	152
153	Chapter LXXXXXXXV	153
154	Chapter LXXXXXXXVI	154
155	Chapter LXXXXXXXVII	155
156	Chapter LXXXXXXXVIII	156
157	Chapter LXXXXXXXIX	157
158	Chapter LXXXXXXXI	158
159	Chapter LXXXXXXXII	159
160	Chapter LXXXXXXXIII	160
161	Chapter LXXXXXXXIV	161
162	Chapter LXXXXXXXV	162
163	Chapter LXXXXXXXVI	163
164	Chapter LXXXXXXXVII	164
165	Chapter LXXXXXXXVIII	165
166	Chapter LXXXXXXXIX	166
167	Chapter LXXXXXXXI	167
168	Chapter LXXXXXXXII	168
169	Chapter LXXXXXXXIII	169
170	Chapter LXXXXXXXIV	170
171	Chapter LXXXXXXXV	171
172	Chapter LXXXXXXXVI	172
173	Chapter LXXXXXXXVII	173
174	Chapter LXXXXXXXVIII	174
175	Chapter LXXXXXXXIX	175
176	Chapter LXXXXXXXI	176
177	Chapter LXXXXXXXII	177
178	Chapter LXXXXXXXIII	178
179	Chapter LXXXXXXXIV	179
180	Chapter LXXXXXXXV	180
181	Chapter LXXXXXXXVI	181
182	Chapter LXXXXXXXVII	182
183	Chapter LXXXXXXXVIII	183
184	Chapter LXXXXXXXIX	184
185	Chapter LXXXXXXXI	185
186	Chapter LXXXXXXXII	186
187	Chapter LXXXXXXXIII	187
188	Chapter LXXXXXXXIV	188
189	Chapter LXXXXXXXV	189
190	Chapter LXXXXXXXVI	190
191	Chapter LXXXXXXXVII	191
192	Chapter LXXXXXXXVIII	192
193	Chapter LXXXXXXXIX	193
194	Chapter LXXXXXXXI	194
195	Chapter LXXXXXXXII	195
196	Chapter LXXXXXXXIII	196
197	Chapter LXXXXXXXIV	197
198	Chapter LXXXXXXXV	198
199	Chapter LXXXXXXXVI	199
200	Chapter LXXXXXXXVII	200

LIST OF FIGURES

Figure 1. Log of retention time vs. carbon number of fatty acids ----- Appendix i

THE UNIVERSITY OF CHICAGO

THE UNIVERSITY OF CHICAGO LIBRARY

1950

1950

1950

1950

1950

1950

1950

1950

1950

1950

1950

1950

1950

1950

1950

1950

1950

1950

1950

1950

1950

1950

1950

1950

1950

INTRODUCTION

Rapeseed is an important crop in Western Canada. Production of 22.8 million bushels of rapeseed in 1965 was almost double the 1964 production of 13.2 million bushels. The 1966 crop is estimated to be similar to that of 1965. Almost 50% of the crop is exported as whole rapeseed. The remainder is chiefly for production of edible oils for domestic consumption.

Rapeseed meal is a high protein by-product of oil extraction and a potential substitute for soybean meal and other vegetable protein meals in livestock and poultry diets. Its use in feeds, especially starting and breeding swine diets, has been limited by regulation. Recent changes in processing methods have made rapeseed meal more attractive as a protein supplement and have resulted in removal in 1966 of the restrictive ruling of the Feeds Act.

Two species of rapeseed are grown in Western Canada: Brassica napus L., commonly called Argentine rapeseed and Brassica campestris L., commonly called Polish rapeseed. The Polish type usually has a lower protein and oxazolidinethione content than the Argentine type. Oxazolidinethione and other compounds in rapeseed inhibit thyroid function, and therefore may have adverse effects on growth.

Research information is limited on the use of rapeseed meal in swine reproduction diets. Therefore, studies were undertaken to determine the effects of including solvent-extracted rapeseed meal in swine diets on growth of female breeding stock, on their reproductive and lactational performance and on the performance of their offspring.

THE PROBLEM

The first part of the paper is devoted to a discussion of the problem of the existence of a solution of the system of equations (1) for a given set of initial conditions. It is shown that the system of equations (1) has a unique solution for a given set of initial conditions if the functions $f_i(x, y, z, t)$ are continuous and satisfy the Lipschitz condition.

In the second part of the paper, the problem of the stability of the solution of the system of equations (1) is considered. It is shown that the solution of the system of equations (1) is stable if the functions $f_i(x, y, z, t)$ are continuous and satisfy the Lipschitz condition. The stability of the solution of the system of equations (1) is also considered for a given set of initial conditions.

The third part of the paper is devoted to a discussion of the problem of the existence of a solution of the system of equations (1) for a given set of initial conditions. It is shown that the system of equations (1) has a unique solution for a given set of initial conditions if the functions $f_i(x, y, z, t)$ are continuous and satisfy the Lipschitz condition.

In the fourth part of the paper, the problem of the stability of the solution of the system of equations (1) is considered. It is shown that the solution of the system of equations (1) is stable if the functions $f_i(x, y, z, t)$ are continuous and satisfy the Lipschitz condition. The stability of the solution of the system of equations (1) is also considered for a given set of initial conditions.

REVIEW OF LITERATURE

Bell (1955), in a review article, stated that published German experiments on the nutritive value of rapeseed meal go back to 1872. He also stated that the first North American report on this subject appeared in 1944. Today the use of rapeseed meal as a protein supplement in swine diets is not as extensive as it might be when quality, availability and price are taken into consideration. Reluctance to use rapeseed meal in some livestock diets stems from reports of adverse effects on growth and reproduction associated with undesirable principles in rapeseed meal. A recent review (Bowland, Clandinin and Wetter, 1965) discusses the nutritive value of rapeseed meal, and shows that a much better quality product is being produced today than 10 to 15 years ago.

A. Goitrogenic Properties of Rapeseed Meal

1. Nature of the potentially toxic factors

Chesney, Clawson and Webster in 1928 first observed the presence of goitrogens in food, when experimental rabbits on a fresh cabbage diet showed up to 200-fold thyroid enlargement. Kennedy and Purves (1941) first reported that rapeseed had goitrogenic properties. Rapeseed and mustard seed were known to be rich in glucosides, and when these seeds were fed to rats, goiters were produced. After feeding rapeseed meal for 3 weeks, thyroid enlargement reached a plateau, although after 9 weeks on rapeseed meal the thyroid re-accumulated some colloid while retaining its hypertrophy. Similar results were obtained recently by Manns (1962) with rats and pigs and by Bowland and Standish (1966) with rats.

In rapeseed meal, thioglucosides are the chief undesirable principles. On enzymatic hydrolysis these glucosides give rise to mustard oils,

CHAPTER 10

THEORY OF THE EARTH AND ITS HISTORY

The theory of the earth and its history is a branch of geology which deals with the origin and development of the earth and its various parts. It is a science which seeks to explain the causes of the various geological phenomena which we observe in nature.

The theory of the earth and its history is a branch of geology which deals with the origin and development of the earth and its various parts. It is a science which seeks to explain the causes of the various geological phenomena which we observe in nature.

THEORY OF THE EARTH AND ITS HISTORY

THEORY OF THE EARTH AND ITS HISTORY

The theory of the earth and its history is a branch of geology which deals with the origin and development of the earth and its various parts. It is a science which seeks to explain the causes of the various geological phenomena which we observe in nature.

The theory of the earth and its history is a branch of geology which deals with the origin and development of the earth and its various parts. It is a science which seeks to explain the causes of the various geological phenomena which we observe in nature.

THEORY OF THE EARTH AND ITS HISTORY

THEORY OF THE EARTH AND ITS HISTORY

THEORY OF THE EARTH AND ITS HISTORY

isothiocyanate and oxazolidinethione. Kjaer, Conti and Jensen (1953) began a series of investigations into the natural mustard oils and their thioglucoside precursors. The first thioglucoside isolated from plant sources was sinigrin from black mustard seed by Bussy in 1840 (cited by Bell and Belzile, 1965), which when treated with the enzyme myrosinase, produced allyl isothiocyanate. Sinalbin is a thioglucoside in rapeseed and gives rise to glucose and p-hydroxybenzyl isothiocyanate upon hydrolysis. Clandinin (1961) ascribes the sharp bitter taste of rapeseed to this isothiocyanate.

Myrosinase is an enzyme found in rapeseed. The enzyme was first isolated by Boutron and Fremy in 1840 (cited by Bell and Belzile, 1965) and called myrosin. Myrosinase may be obtained by extraction of Brassica seeds and the enzyme catalyzes the hydrolysis of mustard oil thioglucosides to isothiocyanates. Kjaer, Conti and Jensen (1953) isolated and identified three volatile mustard oils and found evidence for three major and three minor thioglucosides in rapeseed. Progoitrin and gluconapin are the major non-volatile thioglucosides. Astwood, Greer and Ettlinger (1949) and Carrol (1949) isolated and characterized (-)-5-vinyl-2-oxazolidinethione from rapeseed. This compound was found to be goitrogenic for rats and humans and was given the name goitrin. The glucoside precursor of goitrin was named progoitrin. Progoitrin, under the action of myrosinase (Pitt-Rivers, 1950), gives rise to 3-butenyl isothiocyanate which may give rise to goitrin by cyclization. Clandinin, Renner and Robblee (1959) found that heat increased the goitrogenicity of rapeseed meal by conversion of isothiocyanate to non-volatile goitrin.

2. Action of goitrogens

Rapeseed meal has been shown to be goitrogenic for various non-

the same way as the other two, but the first is the only one that is not a

simple case of the other two, and the second is the only one that is not a

simple case of the other two, and the third is the only one that is not a

simple case of the other two, and the fourth is the only one that is not a

simple case of the other two, and the fifth is the only one that is not a

simple case of the other two, and the sixth is the only one that is not a

simple case of the other two, and the seventh is the only one that is not a

simple case of the other two, and the eighth is the only one that is not a

simple case of the other two, and the ninth is the only one that is not a

simple case of the other two, and the tenth is the only one that is not a

simple case of the other two, and the eleventh is the only one that is not a

simple case of the other two, and the twelfth is the only one that is not a

simple case of the other two, and the thirteenth is the only one that is not a

simple case of the other two, and the fourteenth is the only one that is not a

simple case of the other two, and the fifteenth is the only one that is not a

simple case of the other two, and the sixteenth is the only one that is not a

simple case of the other two, and the seventeenth is the only one that is not a

simple case of the other two, and the eighteenth is the only one that is not a

simple case of the other two, and the nineteenth is the only one that is not a

simple case of the other two, and the twentieth is the only one that is not a

simple case of the other two, and the twenty-first is the only one that is not a

simple case of the other two, and the twenty-second is the only one that is not a

simple case of the other two, and the twenty-third is the only one that is not a

simple case of the other two, and the twenty-fourth is the only one that is not a

simple case of the other two, and the twenty-fifth is the only one that is not a

simple case of the other two, and the twenty-sixth is the only one that is not a

simple case of the other two, and the twenty-seventh is the only one that is not a

simple case of the other two, and the twenty-eighth is the only one that is not a

simple case of the other two, and the twenty-ninth is the only one that is not a

simple case of the other two, and the thirtieth is the only one that is not a

ruminants. Pettit et al (1944) reported that 20% rapeseed meal in a chick starter diet caused thyroid hypertrophy. Clandinin and Bayly (1960) reported that 15% Polish or 5% Argentine type rapeseed meal doubled thyroid-to-body-weight ratios in chicks. Blakely and Anderson (1948a), feeding 20% rapeseed meal to turkey poults, demonstrated a five to six-fold increase in thyroid weight. Hussar and Bowland (1959a) and Manns et al (1963) observed reduced standard metabolic rates in rats fed 10% rapeseed meal. These two authors also showed that in swine, serum protein bound iodine (PBI) was reduced when rapeseed meal was fed from weaning to market weight. They demonstrated thyroid hypertrophy due to feeding of rapeseed meal. There is little evidence to show that rapeseed meal is goitrogenic for ruminants. For example, Bezeau et al (1960) reported no thyroid enlargement occurred in ewes fed rapeseed meal as 30% of their diet.

The feeding of iodide has been shown to partially correct goitrogenic effects of rapeseed meal (Bell and Belzile, 1965, Kratzer et al, 1954, Purves, 1943). Iodinated casein or thyroxine feeding have helped prevent goiters to some degree (Blakely and Anderson, 1948b, Kratzer et al, 1954, Purves, 1943). Kennedy and Purves (1941) showed that hypophysectomy prevented thyroid hypertrophy. Thyroxine, but not iodine or diiodotyrosine prevented hyperplasia due to rapeseed feeding in rats. Therefore, it appears that rapeseed goitrogens interfere with thyroxine synthesis, and that the anterior pituitary, producing thyroid stimulating hormone (TSH), causes thyroid hypertrophy and hyperplasia. Greer (1962) classified antithyroid compounds according to their modes of action. Oxazolidine-thione, apparently interferes with the organic binding of iodine, in which the most sensitive reaction is the coupling of two iodinated tyrosine molecules to form thyroxine or triiodothyronine, which are the thyroid

hormones. Iodination of tyrosine may also be impaired. Isothiocyanates that have significant antithyroid activities include only those capable of cyclizing to form oxazolidinethione. Greer (1962) stated that few antithyroid substances are capable of blocking thyroxine action once it has left the thyroid gland.

B. Nutritive Value of Rapeseed Meal

1. Effect of processing methods

Clandinin, Renner, and Robblee (1959) showed that high temperature processing of rapeseed meal in the expeller process led to a 25% reduction in lysine content of the protein of rapeseed meal. Clandinin and Tajcnar (1961) reported that the lysine content of expeller-processed meal varied from 3.69 to 5.37% of the protein and lysine level was positively correlated with percent fat, which varied from 5.2 to 9.7%. Less than 6% residual fat in expeller-processed meal was accompanied by a marked reduction of lysine content. Clandinin (1965) found an average of 5.5% lysine in protein of fifteen meals from prepress plus solvent process plants. Finlayson (1965) reported a lysine content of 6.6% of the protein in meal from a straight solvent-extraction plant.

Processing of rapeseed meal affects thioglucoside content by activation or deactivation of myrosinase and by chemical modification of thioglucosides. High temperatures and high moisture content lead to rapid hydrolysis of thioglucosides to isothiocyanates (Mustakas et al, 1962). Only solvent or prepress plus solvent extraction processes are used in Canada now, with the exception of one unit in Eastern Canada. These processes result in a meal which is myrosinase-free (Bell and Belzile, 1965) and which contains unhydrolyzed thioglucosides. Clandinin (1965) tested ten different meals at levels of 15% in chick diets. He reported

that chicks fed diets containing prepress-solvent and solvent-extracted rapeseed meals grew 94% and 94.5%, respectively, as rapidly as chicks on a soybean meal control diet.

2. Value in starting diets

In starting swine diets, acceptability and palatability of a diet are important to ensure adequate intake of feed. The sharp bitter taste of rapeseed meal (Clandinin, 1961) may result in decreased feed intake. Bowland (1957) observed that pigs did not accept a prestarter diet containing 2 to 10% rapeseed meal (expeller-extracted) when alternative diets were available. Manns (1962) observed no depression in feed intake when all the soybean meal in a 13% soybean meal diet was replaced isonitrogenously by Polish solvent-extracted rapeseed meal. Seale in 1952 (cited by Bowland, 1965a) showed that 13 to 26% rapeseed meal in diets for pigs from 18 to 36 kg body weight did not adversely affect feed consumption. Stothers in 1964, (cited by Bowland, 1965a) observed a depression of 20% in feed intake when rapeseed meal replaced all the soybean meal (10.4%) in diets for 14 kg pigs. In general, levels of 10% or less of rapeseed meal in pig starting diets appear not to affect intake.

Hussar and Bowland (1959a) observed that average daily gain was not significantly depressed in group-fed pigs from 6 to 16 kg liveweight when they were fed 2% or 10% expeller-extracted rapeseed meal in substitution for soybean meal. In individually-fed pigs, a 0.06 kg per day depression in gain was observed, although efficiency of feed utilization was not affected. Manns and Bowland (1963a) reported that if 25% of the soybean meal were replaced with solvent-extracted rapeseed meal, gain was not depressed, but at a substitution rate of 50% and 100%, gain

was depressed 0.07 and 0.10 kg per day, respectively, for group-fed pigs from 9 to 23 kg liveweight. Seale in 1952 (cited by Bowland, 1965a) showed that gain and feed efficiency were not affected by inclusion of rapeseed meal in starting diets. Stothers, in 1964 (cited by Bowland, 1965a), later showed that substitution in the diet of starting pigs of 50 or 100% of the soybean meal with rapeseed meal led to a reduced rate of gain but feed efficiency was not affected. In general, levels of 5% or less of rapeseed meal do not appear to affect gain and efficiency of feed utilization of pigs from three weeks of age to 25 kg liveweight, and levels of up to 10% may be acceptable.

3. Value in growing and finishing diets

Frölich and Haring (1937a) fed up to 200gm daily of rapeseed meal to young growing pigs with satisfactory results, but they found that average daily gain was depressed in finishing pigs if more than 150 gm with low mustard oil content (0.11 to 0.13%) and more than 100 gm with high mustard oil content was fed. Cook (1941) observed that substitution of rapeseed meal for meat meal gave poor results with pigs. Seale in 1952 (cited by Bowland, 1965a) compared linseed meal and rapeseed meal in diets for swine from 36 kg to 91 kg liveweight. Pigs fed rapeseed meal made slower and less efficient gains although performance was not affected up to 36 kg liveweight by rapeseed meal. When rapeseed meal replaced only 50% of the linseed meal, gain and feed to gain ratios were comparable to those of pigs on linseed meal diets. Pigs fed rapeseed meal tended to waste feed and consumed feed more slowly than pigs fed linseed meal. Further studies by Seale in 1953 (cited by Bowland, 1965a) showed that in a 35% protein supplement containing 20% meat meal, 20% alfalfa meal and 60% linseed meal, rapeseed meal could replace

all the linseed meal without significantly affecting gain, feed intake and efficiency of feed utilization. Clausen, in Denmark (cited by Bowland, 1965a) observed a marked depression in feed consumption, rate of gain and feed efficiency when rapeseed meal was substituted for milk protein in the diets of 24 pigs started at 24 kg liveweight. Fevrier, in 1957 (cited by Bowland, 1965a) reported on experiments in which 10% to 25% rapeseed meal treated by the André process, involving heat and steam extraction, was fed to swine. Daily gain was lower on the highest level of rapeseed meal. Further steam treatment resulted in improved feed intake, average daily gain and feed efficiency. Hussar and Bowland (1959a) showed that 10%, but not 2%, expeller-extracted Argentine rapeseed meal depressed rate of gain and in some cases depressed feed utilization in growing and market pigs. Manns and Bowland (1963a), replacing 0, 25, 50 and 100% of the soybean meal in diets with rapeseed meal (levels up to 15.6% in growing and up 9.6% in finishing diets), reported that the 25% level did not influence average daily gain, average feed intake or feed efficiency, but at the 50 and 100% levels gain and feed efficiency were depressed in group-fed pigs from 23 to 50 kg liveweight. The addition of lysine at the high level of rapeseed meal feeding had no effect on gain and depressed feed efficiency. In general it appears that levels from 6 to 10% of solvent-extracted rapeseed meal are satisfactory in swine diets from 25 to 90 kg liveweight.

Rapeseed meal appears to have no specific effects on swine carcass lean and fat characteristics. Seale in 1952 (cited by Bowland, 1965a) observed improved carcass grades in swine fed rapeseed meal, but he attributed this to slower gains in the finishing period. Later, Seale showed that rapeseed meal had no effect on Record of Performance (R.O.P.)

The first of these is the fact that the
 second of these is the fact that the
 third of these is the fact that the
 fourth of these is the fact that the
 fifth of these is the fact that the
 sixth of these is the fact that the
 seventh of these is the fact that the
 eighth of these is the fact that the
 ninth of these is the fact that the
 tenth of these is the fact that the
 eleventh of these is the fact that the
 twelfth of these is the fact that the
 thirteenth of these is the fact that the
 fourteenth of these is the fact that the
 fifteenth of these is the fact that the
 sixteenth of these is the fact that the
 seventeenth of these is the fact that the
 eighteenth of these is the fact that the
 nineteenth of these is the fact that the
 twentieth of these is the fact that the
 twenty-first of these is the fact that the
 twenty-second of these is the fact that the
 twenty-third of these is the fact that the
 twenty-fourth of these is the fact that the
 twenty-fifth of these is the fact that the
 twenty-sixth of these is the fact that the
 twenty-seventh of these is the fact that the
 twenty-eighth of these is the fact that the
 twenty-ninth of these is the fact that the
 thirtieth of these is the fact that the
 thirty-first of these is the fact that the
 thirty-second of these is the fact that the
 thirty-third of these is the fact that the
 thirty-fourth of these is the fact that the
 thirty-fifth of these is the fact that the
 thirty-sixth of these is the fact that the
 thirty-seventh of these is the fact that the
 thirty-eighth of these is the fact that the
 thirty-ninth of these is the fact that the
 fortieth of these is the fact that the
 forty-first of these is the fact that the
 forty-second of these is the fact that the
 forty-third of these is the fact that the
 forty-fourth of these is the fact that the
 forty-fifth of these is the fact that the
 forty-sixth of these is the fact that the
 forty-seventh of these is the fact that the
 forty-eighth of these is the fact that the
 forty-ninth of these is the fact that the
 fiftieth of these is the fact that the
 fifty-first of these is the fact that the
 fifty-second of these is the fact that the
 fifty-third of these is the fact that the
 fifty-fourth of these is the fact that the
 fifty-fifth of these is the fact that the
 fifty-sixth of these is the fact that the
 fifty-seventh of these is the fact that the
 fifty-eighth of these is the fact that the
 fifty-ninth of these is the fact that the
 sixtieth of these is the fact that the
 sixty-first of these is the fact that the
 sixty-second of these is the fact that the
 sixty-third of these is the fact that the
 sixty-fourth of these is the fact that the
 sixty-fifth of these is the fact that the
 sixty-sixth of these is the fact that the
 sixty-seventh of these is the fact that the
 sixty-eighth of these is the fact that the
 sixty-ninth of these is the fact that the
 seventieth of these is the fact that the
 seventy-first of these is the fact that the
 seventy-second of these is the fact that the
 seventy-third of these is the fact that the
 seventy-fourth of these is the fact that the
 seventy-fifth of these is the fact that the
 seventy-sixth of these is the fact that the
 seventy-seventh of these is the fact that the
 seventy-eighth of these is the fact that the
 seventy-ninth of these is the fact that the
 eightieth of these is the fact that the
 eighty-first of these is the fact that the
 eighty-second of these is the fact that the
 eighty-third of these is the fact that the
 eighty-fourth of these is the fact that the
 eighty-fifth of these is the fact that the
 eighty-sixth of these is the fact that the
 eighty-seventh of these is the fact that the
 eighty-eighth of these is the fact that the
 eighty-ninth of these is the fact that the
 ninetieth of these is the fact that the
 ninety-first of these is the fact that the
 ninety-second of these is the fact that the
 ninety-third of these is the fact that the
 ninety-fourth of these is the fact that the
 ninety-fifth of these is the fact that the
 ninety-sixth of these is the fact that the
 ninety-seventh of these is the fact that the
 ninety-eighth of these is the fact that the
 ninety-ninth of these is the fact that the
 hundredth of these is the fact that the

carcass scores and measurements or on carcass grades. Hussar and Bowland (1959a) observed a significant reduction in loin eye area of pigs on a diet containing 10% rapeseed meal when the pigs were fed individually, but not when group-fed. Carcasses of pigs receiving rapeseed meal appeared to be shorter. Bowland (1951) reported shorter carcasses were obtained when pigs were fed thiouracil (a goitrogen with antithyroid action similar to that of rapeseed goitrogens). Manns and Bowland (1963a) reported no influence on carcass measurements or carcass grades when solvent-extracted rapeseed meal replaced all the soybean meal in the diet. Bell and Belzile (1965) showed that rapeseed meal had little influence on carcass quality, except that it tended to increase loin eye area. Bell, however, showed that feeding of rapeseed screenings to pigs receiving rapeseed meal tended to depress loin eye area, which agrees with the observations of Hussar (1958). Rapeseed screenings contain myrosinase, the enzyme which liberates rapeseed meal goitrogens, and the rapeseed meal used by Bell was reported to be myrosinase-free. In general, however, it has been shown that rapeseed meal has little effect on carcass measurements.

C. Reproduction and lactation

Reports on the effects of rapeseed meal on reproduction and lactation in swine are inconsistent. Frölich and Haring (1937b) stated that up to 400 gm of rapeseed meal per head daily was satisfactory for nursing sows, and 200 gm daily was a maximum for young sows over 60 kg liveweight. Bell, in 1958 (cited by Bell and Belzile, 1965), replacing linseed meal with 7% rapeseed meal in diets for gilts, obtained unsatisfactory reproduction and lactation results. Rapeseed meal at this level depressed number of pigs born alive, and birth weight and weaning weight of litters, especially when Polish type meal was fed. Gilts on rapeseed meal diets lost more

weight from prior to farrowing to the end of lactation than control gilts.

Manns and Bowland (1963a) fed rapeseed meal in gestation and lactation diets to gilts at 0, 3, 6 and 12% of the diet. At 6 and 12% rapeseed meal, the gilts required 2 to 2.5 estrous cycle on the average before they conceived. Addition of lysine to the diet was of no benefit. The same authors reported poor lactation and reproductive performance with rats fed rapeseed meal. Kennedy and Purves (1941) stated that rapeseed meal caused a delay in ovary development. Delayed ovarian development might explain the slow conception observed by Manns. In the experiment of Manns (1962) gilts fed 12% rapeseed meal farrowed numbers of live pigs similar to the control gilts, but birth weight and five week weaning weights were lower for offspring of gilts fed 12% rapeseed meal. Such results suggest lactational inadequacy associated with rapeseed meal feeding. The author suggested that rapeseed meal should be restricted to 3% in pre-gestation, gestation and lactation diets.

D. Composition of Sows' Colostrum and Milk

1. Gross composition

Much work has been done on the yield and composition of sows' colostrum and milk because of the economic importance of early weaning. Sows' milk composition was first studied in the nineteenth century. Von Gohren, in 1865 (cited by Bowland, 1965b) obtained the following percentage values for total solids, fat, protein, milk sugar and ash, respectively: in colostrum--29.87, 9.53, 15.56, 3.84, 0.85, and in milk--13.87, 1.48, 7.91, 2.65, and 1.14.

In a review, Bowland (1965b) cited a review of von Neuhaus (1961) and, in a summary of data appearing in the literature since 1930, obtained the following mean values for sows' milk; percent total solids, fat,

protein, lactose, and ash respectively: 19.4, 7.2, 6.1, 4.8 and 0.96. Other studies of gross composition of sows' milk are those of Braude et al, (1947), who first used oxytocin for milk letdown in sows; Bowland et al (1949a and b), who also studied levels of vitamins A, C and D; Smith (1959a and b) and Evans (1959).

Perrin (1955) showed that as lactation progressed, percent total solids fell rapidly from about 30% at parturition to about 20% at 12 hours postpartum, and was approximately 21% from the second to the eighth week of lactation. Percent fat averaged 7 to 8% for the first day and increased to 9 to 10% for the duration of lactation. Protein was high (near 20%) at parturition and fell in the following 24 hours to 6 to 7%, where it stayed for the rest of lactation. Lactose levels went from 2.5% at parturition to 5% during the second week of lactation. Percent ash increased from 0.6 to 1.0%. Vitamin A (Bowland, 1965b) varied from 145 μ g per 100 ml in colostrum to a low of 11 μ g per 100 ml in sows' milk. Perrin (1955) felt that 4 to 7 days were required for the transition of colostrum to milk.

Beacom and Bowland (1951) reported essential amino acid levels in sows' colostrum and milk. Colostrum, compared to milk, had higher levels of threonine, phenylalanine, valine and leucine and lower levels of lysine and methionine expressed as a percent of the protein. All the amino acids measured except histidine were lower in sows' milk protein than in cows' milk protein.

Rook (1961) reviewed the causes of variation in chemical composition of milk of the cow. He noted effects of season, breed and age of animals, lactational stage and feeding regimen. In the cow, during a milking period, the first milk may have levels of 1% fat and strippings may have levels of 10% fat. In contrast, Whittlestone (1952) has noted that in the

The first part of the report deals with the general situation of the country and the progress of the work. It is followed by a detailed account of the various projects and the results achieved. The report concludes with a summary of the work done and the prospects for the future.

The second part of the report deals with the financial situation of the country. It gives a detailed account of the various sources of revenue and the expenditure incurred. It also gives a summary of the financial results for the year and the prospects for the future. The third part of the report deals with the administrative situation of the country. It gives a detailed account of the various departments and the work done by them. It also gives a summary of the administrative results for the year and the prospects for the future.

The fourth part of the report deals with the social situation of the country. It gives a detailed account of the various social problems and the work done to solve them. It also gives a summary of the social results for the year and the prospects for the future. The fifth part of the report deals with the educational situation of the country. It gives a detailed account of the various educational institutions and the work done by them. It also gives a summary of the educational results for the year and the prospects for the future.

sow, fat remains dispersed and fat content is constant throughout a milking. In cows (Rook, 1961), thyroid hormone leads to an increase in milk yield, a variable increase in percent fat and a small increase in lactose. Thiouracil, a goitrogen, has reverse effects.

The effects of feeding on sow milk composition have not been studied extensively. Smith, (1959b) observed that percent fat increased, percent protein showed no change and percent lactose decreased with decreasing feed intake of sows. Jarl, in 1951 (cited by Altschul, 1958), showed that rapeseed meal in sow diets depressed percent fat in the milk. Willet and Maruyama (1946) reported that the level of fat in sows' milk increased from 6.1% on a diet containing 2.8% fat to 9.6% on a diet containing 27.1% fat. DeMan and Bowland (1963) found that the inclusion of 15% stabilized tallow in sow diets resulted in levels of 8.4% fat in the milk compared to 6.2% fat in milk of sows on standard diets. Bowland et al (1951) observed that pasture-fed sows had significantly lower fat levels in their milk and lower protein levels in first day colostrum than dry-lot fed sows.

2. Fatty acid composition of milk fat

De la Mare and Shorland (1944) demonstrated by ester fractionation methods that the fat of hand-milked Berkshire sows contained 2.4, 1.8, 20.3, 6.1, 8.8, 35.0, 14.0 and 3.6 moles per cent, respectively of steam volatile fatty acids, myristic acid, palmitic acid, stearic acid, hexadecenoic acid, oleic acid, octadecadienoic acid, and 20 to 22-carbon acids. In a recent study, DeMan and Bowland (1963) reported levels of fatty acids as a percent of milk fat of sows fed a diet to meet U.S. N.R.C. requirements were as follows: oleic, 35.3; palmitic, 30.3; linoleic, 13.0; palmitoleic, 9.9; stearic, 4.0; myristic, 3.3; linolenic, 2.5; lauric, 0.3;

and capric, 0.2. The corresponding composition of colostrum fat was: oleic, 41.7; palmitic, 22.5; linoleic, 20.9; palmitoleic, 5.0; stearic, 5.7; myristic, 1.4; and linolenic, 2.4. Salmon-Legagneur (1964) fed sows 8% levels of maize and coconut oils during pregnancy and lactation. Groups fed maize oil during pregnancy and no supplemental oil during lactation (M-A) and groups fed maize oil during pregnancy and coconut oil during lactation (M-C) had high levels of linoleic acid in milk fat, and M-C groups had more lauric acid in milk fat than controls. Adding oil to diets led to increasing proportions of fatty acids in milk resulting from dietary sources and decreasing proportions from endogenous sources. Tollerz and Lindberg (1965) showed that addition of 6.3% cottonseed oil to sow diets led to doubling of linoleic acid content of milk fat, and 6.3% linseed oil (58% linolenic acid) raised linolenic acid content of milk fat from 1.1 to 18.3%. A compensating decrease in 14 and 16-carbon fatty acids was observed. Milk fat adjusted to dietary fat patterns in two to three days. A level of 16.7% cottonseed meal in diets led to 27% levels of linoleic acid in milk fat of sows. Garton (1963) stated that in fasting cows, levels of 4 to 14-carbon fatty acids were decreased in milk fat, and suggested that rapid synthesis of fat leads to increasing proportions of short chain fatty acids. However, his report suggested that in sows' milk fat the opposite appears to be true.

Ashworth, Ramaiah, and Keyes (1966) compared the fatty acid composition of milk from seven species of animals. They observed that short chain fatty acids (below 16-carbon) were much lower in milk fat of non-ruminants than in that of the ruminant animals listed.

DeMan and Bowland (1963) demonstrated higher levels of capric, lauric, myristic, palmitic and palmitoleic acids and lower levels of 18-carbon acids in sows' milk than in colostrum. Stinson (1966) reported, in

the first of these is the fact that the
 second of these is the fact that the
 third of these is the fact that the
 fourth of these is the fact that the
 fifth of these is the fact that the
 sixth of these is the fact that the
 seventh of these is the fact that the
 eighth of these is the fact that the
 ninth of these is the fact that the
 tenth of these is the fact that the
 eleventh of these is the fact that the
 twelfth of these is the fact that the
 thirteenth of these is the fact that the
 fourteenth of these is the fact that the
 fifteenth of these is the fact that the
 sixteenth of these is the fact that the
 seventeenth of these is the fact that the
 eighteenth of these is the fact that the
 nineteenth of these is the fact that the
 twentieth of these is the fact that the
 twenty-first of these is the fact that the
 twenty-second of these is the fact that the
 twenty-third of these is the fact that the
 twenty-fourth of these is the fact that the
 twenty-fifth of these is the fact that the
 twenty-sixth of these is the fact that the
 twenty-seventh of these is the fact that the
 twenty-eighth of these is the fact that the
 twenty-ninth of these is the fact that the
 thirtieth of these is the fact that the
 thirty-first of these is the fact that the
 thirty-second of these is the fact that the
 thirty-third of these is the fact that the
 thirty-fourth of these is the fact that the
 thirty-fifth of these is the fact that the
 thirty-sixth of these is the fact that the
 thirty-seventh of these is the fact that the
 thirty-eighth of these is the fact that the
 thirty-ninth of these is the fact that the
 fortieth of these is the fact that the
 forty-first of these is the fact that the
 forty-second of these is the fact that the
 forty-third of these is the fact that the
 forty-fourth of these is the fact that the
 forty-fifth of these is the fact that the
 forty-sixth of these is the fact that the
 forty-seventh of these is the fact that the
 forty-eighth of these is the fact that the
 forty-ninth of these is the fact that the
 fiftieth of these is the fact that the
 fifty-first of these is the fact that the
 fifty-second of these is the fact that the
 fifty-third of these is the fact that the
 fifty-fourth of these is the fact that the
 fifty-fifth of these is the fact that the
 fifty-sixth of these is the fact that the
 fifty-seventh of these is the fact that the
 fifty-eighth of these is the fact that the
 fifty-ninth of these is the fact that the
 sixtieth of these is the fact that the
 sixty-first of these is the fact that the
 sixty-second of these is the fact that the
 sixty-third of these is the fact that the
 sixty-fourth of these is the fact that the
 sixty-fifth of these is the fact that the
 sixty-sixth of these is the fact that the
 sixty-seventh of these is the fact that the
 sixty-eighth of these is the fact that the
 sixty-ninth of these is the fact that the
 seventieth of these is the fact that the
 seventy-first of these is the fact that the
 seventy-second of these is the fact that the
 seventy-third of these is the fact that the
 seventy-fourth of these is the fact that the
 seventy-fifth of these is the fact that the
 seventy-sixth of these is the fact that the
 seventy-seventh of these is the fact that the
 seventy-eighth of these is the fact that the
 seventy-ninth of these is the fact that the
 eightieth of these is the fact that the
 eighty-first of these is the fact that the
 eighty-second of these is the fact that the
 eighty-third of these is the fact that the
 eighty-fourth of these is the fact that the
 eighty-fifth of these is the fact that the
 eighty-sixth of these is the fact that the
 eighty-seventh of these is the fact that the
 eighty-eighth of these is the fact that the
 eighty-ninth of these is the fact that the
 ninetieth of these is the fact that the
 ninety-first of these is the fact that the
 ninety-second of these is the fact that the
 ninety-third of these is the fact that the
 ninety-fourth of these is the fact that the
 ninety-fifth of these is the fact that the
 ninety-sixth of these is the fact that the
 ninety-seventh of these is the fact that the
 ninety-eighth of these is the fact that the
 ninety-ninth of these is the fact that the
 hundredth of these is the fact that the

agreement with DeMan and Bowland, that myristic, palmitic, palmitoleic, stearic, oleic, linoleic, arachidic and gadoleic acids were respectively, 1.8, 22.8, 4.3, 4.5, 35.4, 27.3, 3.1 and 0.5% of colostrum fat of sows, and 3.9, 30.8, 11.4, 3.8, 32.7, 14.3, 1.7 and 0.3% respectively of sows' milk fat. Fatty acids up to palmitic were significantly lower in colostrum fat and fatty acids from stearic to the 20-carbon fatty acids were lower in milk fat.

Duncan and Garton (1966), in a recent study of sows' colostrum and milk fat, demonstrated a reciprocal relationship between the levels of C 16:0 and C 18:1 fatty acids. They also demonstrated that C 16:0 (palmitic acid) was preferentially esterified to the second carbon atom of the glycerol portion of triglycerides, which observation agrees with observations of Stinson (1966).

EXPERIMENTAL

A. Objectives

These experiments were designed to study the effects on growth, gestation and lactation when 8% solvent-extracted rapeseed meal was substituted on a protein equivalent basis for soybean meal in the diets.

Specific experiments were designed to study:

- (1) a) Rate of gain and efficiency of feed utilization of gilts through growing and pre-gestation periods,
b) reproductive performance of sows for two reproductive cycles,
c) gross composition of colostrum and milk and fatty acid composition of colostrum and milk fat,
- (2) a) rate of gain and efficiency of feed utilization from weaning to market weight of first litter progeny from sows,
b) digestible energy and nitrogen, and metabolizable energy and nitrogen of the diets when the progeny were 6 weeks of age and also when they were 40 kg liveweight.

B. Methods and Procedures

1. General

Experiment 1 was conducted from September, 1965 to December, 1966 and experiment 2, using progeny from sows in experiment 1, continued from July, 1966 to December, 1966.

Experiment 1 dealt with growth and reproductive performance of 24 gilts through two reproductive cycles. Feed consumption, daily gain, and efficiency of feed utilization were measured during growth and up to the first farrowing. Sows' colostrum and milk composition was studied after

Appendix

Page 10

The following table shows the results of the survey conducted in the year 2000. The data was collected from 100 respondents and is presented in the following table.

The following table shows the results of the survey conducted in the year 2000.

The following table shows the results of the survey conducted in the year 2000.

The following table shows the results of the survey conducted in the year 2000.

The following table shows the results of the survey conducted in the year 2000.

Table 1

The following table shows the results of the survey conducted in the year 2000.

The following table shows the results of the survey conducted in the year 2000.

The following table shows the results of the survey conducted in the year 2000.

The following table shows the results of the survey conducted in the year 2000.

The following table shows the results of the survey conducted in the year 2000.

Table 2

The following table shows the results of the survey conducted in the year 2000.

The following table shows the results of the survey conducted in the year 2000.

The following table shows the results of the survey conducted in the year 2000.

Table 3

Table 4

The following table shows the results of the survey conducted in the year 2000.

The following table shows the results of the survey conducted in the year 2000.

Table 5

The following table shows the results of the survey conducted in the year 2000.

The following table shows the results of the survey conducted in the year 2000.

The following table shows the results of the survey conducted in the year 2000.

The following table shows the results of the survey conducted in the year 2000.

the first farrowing with milk samples being obtained from 16 sows.

Experiment 2 was conducted with 32 first litter offspring of sows in experiment 1. Feed consumption, daily gain, efficiency of feed utilization, digestibility of nitrogen and energy, and carcass characteristics were studied. The pigs were identified at birth by ear notching and routinely treated for anemia prevention by injection of 2 ml of Pigdex 100 (contains 100 mg of injectable iron per ml) at 4 days of age. Males were castrated following weaning at 21 days of age. Pigs were treated for roundworms (ascarids) with a piperazine derivative and were vaccinated for erysipelas.

2. Formulation of experimental diets and methods of feeding

Diets were formulated to meet requirements as defined in the National Academy of Sciences--National Research Council Nutrient Requirements of Swine (1964; 1967, unpublished). In experiment 2, the starting diets were fed from the age of 3 weeks to 9 weeks; the growing diets from 9 weeks of age to 55 kg liveweight and the finishing diets from 55 kg to market weight of 90 kg. The gilts in experiment 1 were on a standard diet without rapeseed meal up to 12 weeks of age at which time they were allotted to experimental diets as outlined for experiment 2.

At 90 kg liveweight the gilts were put on the reproduction diets and individually fed 2.7 kg daily of feed. Two weeks after breeding and until transferred to farrowing pens, they were restricted to 2.3 kg of feed daily. On the day of farrowing they were given 0.45 kg of feed and 0.45 kg of bran. The bran was eliminated on the fourth day after farrowing. The feed intake of sows was increased to 2.3 kg daily plus 0.3 kg daily per pig in the litter by the seventh day after farrowing. At 20 days after farrowing, feed was reduced to 2.3 kg daily and held at

THE FIRST OF THESE IS THE FACT THAT THE

RESEARCHERS HAVE BEEN ABLE TO

IDENTIFY THE FACTORS WHICH

ARE RESPONSIBLE FOR THE

DIFFERENCES IN THE

REACTIVITY OF THE

DIFFERENT GROUPS

AND THE FACT THAT

THE REACTIVITY OF THE

DIFFERENT GROUPS

IS NOT THE SAME

IN ALL CASES

AND THAT THE

REACTIVITY OF THE

DIFFERENT GROUPS

IS NOT THE SAME

IN ALL CASES

AND THAT THE

REACTIVITY OF THE

DIFFERENT GROUPS

IS NOT THE SAME

IN ALL CASES

AND THAT THE

REACTIVITY OF THE

DIFFERENT GROUPS

this level throughout the second gestation period. The sows, during the second farrowing and lactation were fed similarly to the method used in the first farrowing and lactation. Experiment 1 was terminated 21 days after the second farrowing of each sow at which time the litters were weaned.

Tables 1 and 2 show the formulation and composition of the experimental diets. The rapeseed meal diets were adapted from the basal diets by substituting rapeseed meal for wheat and soybean meal so that the diets remained isonitrogenous and approximately isocaloric. Table 2A lists the composition of the rapeseed meal used in the diets. All diets were ground, mixed and bagged at the University Livestock Farm elevator.

3. Experiment 1

a) Sow growth and reproduction experiments

In experiment 1, 24 gilts (6 littermate groups of 4 gilts each) were assigned to four lots which received the treatments shown in Table 3. One gilt from each of the 6 littermate groups was assigned to each of the four lots. The gilts averaged 29 kg liveweight when allotted. For the growing and finishing periods (29 to 90 kg), the gilts were group-fed in eight pens of three pigs each, and when they reached a weight of 90 kg, they were transferred to six pens of four pigs each fed individually. They were housed in a temperature-controlled building on concrete floors.

The gilts were mated to boars, according to the schedule in Table 4, at an average age of 230 days. The first mating period was for 43 days, from February 2 to March 17 to allow the possibility of two estrous periods, and the second mating period was from weaning of the first litter to 64 days after weaning according to the schedule in Table 4.

Table 1. Formulation of the diets on an air-dry basis

Diet	<u>Starter</u>		<u>Grower</u>		<u>Finisher</u>		<u>Reproduction</u>	
Rapeseed meal added	-	+	-	+	-	+	-	+
Diet number	1	2	3	4	5	6	7	8
Ingredients	%							
Wheat	60.9	59.9	41.0	40.0	32.5	31.5	26.5	25.5
Barley	-	-	41.05	41.05	33.0	33.0	26.95	26.95
Oats	-	-	-	-	20.0	20.0	30.0	30.0
Oat groats	4.10	4.10	-	-	-	-	-	-
Dried skim milk powder	5.00	5.00	-	-	-	-	-	-
Alfalfa meal	-	-	2.00	2.00	3.00	3.00	4.00	4.00
Soybean meal (44%)	10.00	3.00	10.00	3.00	8.00	1.00	9.00	2.00
Rapeseed meal	-	8.00	-	8.00	-	8.00	-	8.00
Fishmeal (72%)	7.00	7.00	2.00	2.00	-	-	-	-
Meat meal (55%)	1.20	1.20	2.00	2.00	2.00	2.00	2.00	2.00
Stabilized fat	3.00	3.00	-	-	-	-	-	-
Sucrose	5.00	5.00	-	-	-	-	-	-
Molasses	2.40	2.40	-	-	-	-	-	-
Ground limestone	0.20	0.20	0.50	0.50	0.50	0.50	0.50	0.50
Dicalcium phosphate	-	-	0.60	0.60	0.20	0.20	0.30	0.30
Iodized salt	0.50	0.50	0.50	0.50	0.50	0.50	0.50	0.50
Zinc sulfate	0.05	0.05	0.05	0.05	0.05	0.05	-	-
Trace mineral mix ¹	0.15	0.15	0.10	0.10	0.10	0.10	0.10	0.10
Vitamin B-complex mix ²	0.30	0.30	0.10	0.10	0.10	0.10	0.10	0.10
Vitamins A and D ₂ ³	+	+	+	+	+	+	+	+
Vitamin B ₁₂	-	-	-	-	-	-	0.05	0.05
Antibiotic supplement								
TM-10 ⁴	0.20	0.20	0.10	0.10	0.05	0.05	-	-

¹Contains the following minerals per kg of mineral mix: cobalt carbonate, 2.28 gm; copper sulphate (CuSO₄·5H₂O), 24.50 gm; ethylene diamine dihydroiodide, 1.30 gm; ferrous carbonate, 234.80 gm; manganous oxide, 47.73 gm; zinc oxide, 2.96 gm; ground limestone, 686.43 gm.

²Contains the following B vitamins per kg of vitamin mix: riboflavin, 4.4 gm; calcium pantothenate, 8.8 gm; niacin, 19.8 gm; choline chloride, 21.4 gm; folic acid, 132.0 mg.

³Diets 1-6: To provide 200,000 I.U. vitamin A and 20,000 I.U. vitamin D₂ per 100 kg.
Diets 7-8: To provide 500,000 I.U. vitamin A and 50,000 I.U. vitamin D₂ per 100 kg.

⁴Contains 22 gm oxytetracycline per kg.

Table 2. Composition of the diets on an air-dry basis

Diet number		1	2	3	4	5	6	7	8
Soybean meal ¹	%	10.0	3.0	10.0	3.0	10.0	3.0	10.0	3.0
Rapeseed meal ¹	%	0.0	8.0	0.0	8.0	0.0	8.0	0.0	8.0
Components									
Gross energy	kcal/kg	4284*	4256*	3943*	3924*	4370 ⁺	4360 ⁺	4220 ⁺	4210 ⁺
Digestible energy (calculated)	kcal/kg	3470	3440	3360	3330	3260	3240	3190	3170
Digestible energy (analysis)	kcal/kg	3644	3609	3254	3174				
Metabolizable energy (analysis)	kcal/kg	3517	3488	3135	3059				
Crude protein (calculated)	%	22.1	22.1	17.5	17.6	15.7	15.8	16.2	16.1
Crude protein (analysis)	%	22.7	21.5	18.0	18.2	16.0	15.7	16.0	16.0
Digestible protein (analysis)	%	19.1	17.7	15.0	15.0				
Metabolizable protein (analysis)	%	12.6	11.2	7.4	7.7				
Lysine (calculated)	%	1.14	1.09	0.84	0.79	0.69	0.64	0.72	0.66
Crude fat (analysis)	%	5.2	5.2	2.2	2.2	2.9	3.2	3.6	3.7
Ash (analysis)	%	3.2	3.4	3.8	4.0	3.3	3.6	4.3	4.0

¹In subsequent tables soybean meal is referred to as SM and rapeseed meal as RM.

*Values obtained by analysis

⁺Calculated values

THE JOURNAL OF THE AMERICAN MEDICAL ASSOCIATION
PUBLISHED WEEKLY

Subscription Rates for 1917							
Country	Single Copy	1 Year	2 Years	3 Years	4 Years	5 Years	6 Years
United States	10c	\$1.00	\$1.80	\$2.50	\$3.20	\$3.90	\$4.60
Canada	10c	\$1.00	\$1.80	\$2.50	\$3.20	\$3.90	\$4.60
Foreign	10c	\$1.00	\$1.80	\$2.50	\$3.20	\$3.90	\$4.60
Advertising Rates							
Per Line per Week	10c						
Per Line per Month	\$2.50						
Per Line per Quarter	\$7.50						
Per Line per Half Year	\$12.50						
Per Line per Year	\$25.00						
Per Column per Week	\$10.00						
Per Column per Month	\$25.00						
Per Column per Quarter	\$75.00						
Per Column per Half Year	\$125.00						
Per Column per Year	\$250.00						
Per Page per Week	\$10.00						
Per Page per Month	\$25.00						
Per Page per Quarter	\$75.00						
Per Page per Half Year	\$125.00						
Per Page per Year	\$250.00						

Published by the American Medical Association, 535 North Dearborn Street, Chicago, Ill.
Subscription Department, 535 North Dearborn Street, Chicago, Ill.
Entered as Second-Class Matter, May 2, 1912, Post Office at Chicago, Ill., under No. 100,000.
Acceptance for mailing at special rate of postage provided for in Act of October 3, 1917, authorized on July 1, 1918.
Postage paid at Chicago, Ill.

Table 2A. Composition of rapeseed meal compared to soybean meal

	Gross energy kcal/kg	DE ¹ kcal/kg	ME ¹ kcal/kg	Crude protein %	Digestible protein %	Lysine %	Crude fat %	Ash %
Rapeseed meal	4380*	3000**	1760 ⁺⁺	34.4*	26.8**	2.0*	1.9*	5.9*
Soybean meal	4200**	3300**	2640 ⁺⁺	45.8**	43.1**	2.9**	1.3**	5.8**

¹DE -- digestible energy

ME -- metabolizable energy

*Values obtained by analysis

+Calculated values

**Values from NRC nutrient requirements of swine (1964)

⁺⁺Values from Clandinin and Robblee (1966)

Table 3. Allotment of gilts to experimental diets -- experiment 1

Lot	1	2	3	4
<u>Dietary treatment</u> ¹				
Growth to 106 kg	SM	RM	RM	SM
106 kg to reproduction	SM	RM	SM	RM
No. of gilts	6	6	6	6
Avg wt. on test kg	28.4	28.4	29.5	29.3
Avg age on test days	84	84	84	84

¹The gilts were group-fed by lots during growth and were individually restricted-fed during reproduction.

THE UNIVERSITY OF CHICAGO LIBRARY

DATE	TIME	FROM	TO	BY	REMARKS
1961	10:00	10:00	10:00	10:00	10:00
1961	10:00	10:00	10:00	10:00	10:00

THE UNIVERSITY OF CHICAGO LIBRARY
 540 EAST 58TH STREET
 CHICAGO, ILL. 60637
 TEL. 777-3000

THE UNIVERSITY OF CHICAGO LIBRARY

DATE	TIME	FROM	TO	BY	REMARKS
1961	10:00	10:00	10:00	10:00	10:00
1961	10:00	10:00	10:00	10:00	10:00

THE UNIVERSITY OF CHICAGO LIBRARY
 540 EAST 58TH STREET
 CHICAGO, ILL. 60637
 TEL. 777-3000

Table 4. Breeding schedule

Gilt litter group number ¹	200	220	240	270	280	290
Breed of gilts ²	Y	LacLandY	YLacY	HYLacY	YLandLacY	YLacY
Boar - first breeding	Y	Y	HY	HY	HY	Lac
Boar - second breeding	Y	Y	Lac	Lac	Y	Lac

¹Each group was 4 littermate gilts allotted across the 4 treatments.

²Y, Lac, Land, and H stand for the breeds Yorkshire, Lacombe, Landrace and Hampshire, respectively.

In both gestation periods, the sows were transferred to farrowing pens 106 days after their last mating, and they were kept there until their litters were weaned at 21 days of age. Weaned pigs were put on a starting diet which contained a level of 0 or 8% rapeseed meal, according to the diet of their dam, until allotted to the growing experiment 2. First litter pigs were weaned throughout June and in early July, 1966, and the sows were transferred immediately to outdoor dry-lot facilities for rebreeding.

b) Lactation experiments

Milk samples were obtained from 16 sows (4 per lot) 8 to 12 hours and 7 days after farrowing of the litters in experiment 1. Pigs were removed from the sow at least 1 hour before milking. Milk letdown was induced by injection of 1 ml of Pitocin (10 IU of oxytocin per ml) into an ear vein. Milk yield was not determined, but a sample of from

THE UNIVERSITY OF CHICAGO

FACULTY				STUDENT		
1900	1901	1902	1903	1904	1905	1906
1907	1908	1909	1910	1911	1912	1913
1914	1915	1916	1917	1918	1919	1920
1921	1922	1923	1924	1925	1926	1927

The University of Chicago is a private research university in Chicago, Illinois. It was founded in 1837 and is one of the oldest and most prominent universities in the United States. The university is known for its commitment to academic excellence and its diverse student body.

The University of Chicago is a private research university in Chicago, Illinois. It was founded in 1837 and is one of the oldest and most prominent universities in the United States. The university is known for its commitment to academic excellence and its diverse student body. The university has a long history of producing world-class scholars and leaders in various fields of study. Its research output is highly influential, and it has a strong reputation for its contributions to the advancement of knowledge.

The University of Chicago is a private research university in Chicago, Illinois. It was founded in 1837 and is one of the oldest and most prominent universities in the United States. The university is known for its commitment to academic excellence and its diverse student body. The university has a long history of producing world-class scholars and leaders in various fields of study. Its research output is highly influential, and it has a strong reputation for its contributions to the advancement of knowledge.

100 to 400 ml was obtained. Milk was collected in amber colored glass containers which were air-tight when sealed. Within 2 hours the sample was placed in a cold storage room at 4 C and within 48 hours the sample was lyophilized in a freeze-drying apparatus.

4. Experiment 2 -- Growth and metabolism experiments

In experiment 2, 4 pigs (2 males and 2 females) from each of 8 litters from lots 1 and 2 of experiment 1 were assigned to 4 lots. The 16 pigs from dams in lot 1 (receiving basal diet) were allotted at random to lots 1 and 4. Pigs from dams in lot 2 (receiving 8% rapeseed meal) were similarly allotted to lots 2 and 3. Male and female pigs were fed separately. Lots 1 and 3 were fed rapeseed meal diets and lots 2 and 4 were fed basal diets (Table 5). The pigs averaged 33 days of age and 5.9 kg liveweight at allotment. They were group-fed in lots of four.

Digestible and metabolizable energy and nitrogen were determined at 6 weeks of age and again at 40 kg liveweight on all male pigs in experiment 2. For the metabolism trials at 6 weeks of age, pigs were housed in small animal cages described by Hussar and Bowland (1959b). The pigs were fed the diets for at least 1 week prior to collection. They were put in the cages within ± 3 days of 6 weeks of age. They were acclimatized to the cages for 3 days, and urine and feces were then collected daily for a 3-day period. Feces were collected on a 12 mesh screen below the cages, and urine was collected

The first part of the paper is devoted to a general discussion of the problem of the existence of solutions of the system of equations (1) for arbitrary values of the parameters α and β . It is shown that the system has solutions for all values of the parameters α and β if the function $f(x)$ is continuous and has a bounded derivative.

2. THE EXISTENCE OF SOLUTIONS OF THE SYSTEM OF EQUATIONS (1)

In the first part of the paper we shall prove the existence of solutions of the system of equations (1) for arbitrary values of the parameters α and β . It is shown that the system has solutions for all values of the parameters α and β if the function $f(x)$ is continuous and has a bounded derivative. The proof is based on the theorem of the existence of solutions of the system of equations (1) for arbitrary values of the parameters α and β if the function $f(x)$ is continuous and has a bounded derivative. The proof is based on the theorem of the existence of solutions of the system of equations (1) for arbitrary values of the parameters α and β if the function $f(x)$ is continuous and has a bounded derivative.

3. THE EXISTENCE OF SOLUTIONS OF THE SYSTEM OF EQUATIONS (1)

In the second part of the paper we shall prove the existence of solutions of the system of equations (1) for arbitrary values of the parameters α and β . It is shown that the system has solutions for all values of the parameters α and β if the function $f(x)$ is continuous and has a bounded derivative. The proof is based on the theorem of the existence of solutions of the system of equations (1) for arbitrary values of the parameters α and β if the function $f(x)$ is continuous and has a bounded derivative. The proof is based on the theorem of the existence of solutions of the system of equations (1) for arbitrary values of the parameters α and β if the function $f(x)$ is continuous and has a bounded derivative.

in pans below the screen and the urine funnelled into approximately 1-liter glass receptacles. The urine was acidified by addition of 25 ml 50% (v/v) H_2SO_4 to the receptacles prior to collection. The total fecal collection and a urine sample were stored at 4 C until placed in a drying oven. The feces were dried in a tray at 60 C in a forced-air oven over a period of 60 hours. The dried feces were allowed to equilibrate with atmospheric moisture for 24 hours. Random samples were shown to contain 8% moisture.

When the pigs reached a liveweight of 40 kg or over on the regular Tuesday weighing, they were transferred to metabolism crates for the second time. These crates were described by Hussar (1958). A 3-day acclimatization period was used and feces and urine were collected separately over a 3-day period. Urine was received in 10-liter plastic pails containing 50 ml 50% H_2SO_4 . Feces and urine were collected, stored, dried and analyzed as for the younger pigs.

the first of these, and which has been the subject of a
 considerable amount of discussion, is the question of
 the "right of the people to know" the truth about
 the activities of the government. This is a question
 which has been raised in many different forms, and
 which has been the subject of much discussion in
 the United States. It is a question which has been
 raised in many different forms, and which has been
 the subject of much discussion in the United States.
 It is a question which has been raised in many
 different forms, and which has been the subject of
 much discussion in the United States. It is a
 question which has been raised in many different
 forms, and which has been the subject of much
 discussion in the United States. It is a question
 which has been raised in many different forms,
 and which has been the subject of much discussion
 in the United States. It is a question which has
 been raised in many different forms, and which
 has been the subject of much discussion in the
 United States. It is a question which has been
 raised in many different forms, and which has
 been the subject of much discussion in the United
 States. It is a question which has been raised in
 many different forms, and which has been the
 subject of much discussion in the United States.

Table 5. Allotment of progeny to experimental diets -- experiment 2

Lot	1	2	3	4
<u>Treatment</u>				
Dams' diet	SM	RM	RM	SM
Diet fed to progeny	SM	RM	SM	RM
No. of pigs	8	8	8	8
Avg wt on test kg	6.0	5.8	5.9	6.0
Avg age on test days	33	31	35	33

5. Methods of chemical analysis

Chemical analyses of feces and urine were similar for both digestion trials except for determination of the energy in urine. The dried feces were ground in a Wiley mill to pass through a 20 mesh screen. Energy in feces and urine was determined by combustion in a Parr oxygen bomb calorimeter. For the determination of energy in urine in the first trials, a 50 ml sample of urine was mixed with 10 gm of soluble starch and dried for 12 hours at 60C in a forced-air oven. The dry material was combusted in the bomb calorimeter and a blank was determined with starch alone. The difference in caloric content between the sample and the blank was taken to represent the energy content of urine. In the trials with 40 kg pigs, a 100 ml sample of urine was lyophilized in a freeze-dryer, and the dry urinary material was combusted directly in the bomb calorimeter. Percent dry matter in urine was determined with 5 ml samples which were lyophilized in small freeze-drying bulbs.

For the determination of nitrogen, the Kjeldahl method was used (AOAC, 1964). To calculate protein levels, a factor of 6.25 was used. Moisture, fat, and ash were determined by the official methods of the

TABLE I			
Summary of the results of the experiments			
Experiment	Time (min)	Distance (m)	Speed (m/s)
1	10	100	10
2	20	200	10
3	30	300	10
4	40	400	10
5	50	500	10
6	60	600	10
7	70	700	10
8	80	800	10
9	90	900	10
10	100	1000	10

The results of the experiments are summarized in Table I. The table shows that the speed of the object is constant at 10 m/s for all experiments. The distance traveled by the object is directly proportional to the time taken for the experiment. The data points are as follows:

Experiment	Time (min)	Distance (m)	Speed (m/s)
1	10	100	10
2	20	200	10
3	30	300	10
4	40	400	10
5	50	500	10
6	60	600	10
7	70	700	10
8	80	800	10
9	90	900	10
10	100	1000	10

The constant speed of 10 m/s is maintained throughout the experiments, indicating a uniform motion. The linear relationship between time and distance is clearly evident from the data presented in Table I.

AOAC (1964).

With colostrum and milk, total solids, protein, ash, solids-not-fat, lactose, and fat as a percent of whole milk, and fatty acids as a percent of fat were determined. Total solids, protein and fat were determined by AOAC (1964) methods, solids-not-fat was determined by difference between percent total solids and percent fat, and lactose was determined by subtracting percent fat, protein and ash from total solids.

To prepare for fatty acid determinations, the freeze-dried colostrum and milk were dissolved in water; and the fat was extracted by the Majonnier method, using a 50/50 (v/v) mixture of ethyl ether and light petroleum ether. The extracted fat was transferred to a 5 ml freeze-drying bulb and esterfied according to the method of DeMan (1964). The esters were injected directly into a gas chromatograph. The instrument used was an F & M model 720 dual column temperature-programmed gas chromatograph equipped with 10 ft (305 cm) columns packed with 20% (w/w) diethyleneglycol succinate (DEGS) on 60/80 mesh firebrick. Column temperature was programmed from 100 to 220 C at a rate of 7.5 per minute. The detector, a thermal conductivity type, was set at 265 C and the injection port at 245 C. Helium carrier gas was used and the flow rate was adjusted to 50 ml per minute. Sample size was 10 to 20 μ l, and the attenuation setting was 32 for the major large peaks, 8 for the smaller peaks and 2 for trace components (below 1%). A Minneapolis-Honeywell strip chart recorder was attached. A planimeter was used to measure peak areas on the chromatograms. To calculate weight percent of the fatty acids, correction factors (DeMan and Bowland, 1963) were applied to area figures and the corrected figures were converted to weight percent by taking them as a percent of total area of all peaks on

the chromatogram. Peaks were identified by 'spiking' esterified fat samples with pure standard fatty acid esters and by a semilogarithmic plot of carbon number vs. logarithm of the retention time (see appendix 1).

6. Methods of statistical analysis

An IBM 7040 computer in the Department of Computing Science was used to statistically analyze all data. A probability of 0.05 was selected as the point of significance for most of the data. For some of the data, particularly that on reproduction, a 10% level of significance was used for interpretation of the results, with some reservations. This level of significance was used because reproductive data of swine tend to vary widely between replicates and within treatments, making statistical analysis difficult. Program BMD02V was used for analysis of variance and program BMD03V was used for analysis of covariance. All data was converted to metric units before statistical analysis. Duncan's new multiple range test (Steele and Torrie, 1960) was used to test differences between multiple means.

Missing data were replaced by the average value of the group from which the data were missing, and in the analysis of variance, error degrees of freedom were reduced by one for each missing value. Data for one sow were missing for the gross and fatty acid composition of milk in experiment 1 and for one pig for the digestibility studies on the growing diets in experiment 2.

The first of these is the fact that the
economy is not in a state of equilibrium
and is therefore subject to fluctuations.

THE PROBLEM OF THE ECONOMY

The first of these is the fact that the

economy is not in a state of equilibrium

and is therefore subject to fluctuations.

The second of these is the fact that the

economy is not in a state of equilibrium

and is therefore subject to fluctuations.

The third of these is the fact that the

economy is not in a state of equilibrium

and is therefore subject to fluctuations.

The fourth of these is the fact that the

economy is not in a state of equilibrium

and is therefore subject to fluctuations.

The fifth of these is the fact that the

economy is not in a state of equilibrium

and is therefore subject to fluctuations.

The sixth of these is the fact that the

economy is not in a state of equilibrium

and is therefore subject to fluctuations.

The seventh of these is the fact that the

RESULTS AND DISCUSSION

A. Growth, Reproduction and Lactation of Sows -- Experiment 1

1. Feed consumption, daily gain and efficiency of feed utilization during pre-gestation

The means for each lot for average daily feed, average daily gain and efficiency of feed utilization for experiment 1 are listed in Table 6. The statistical analysis of these data and treatment means for each factor are given in Table 7.

a) Average daily feed

No significant differences in feed consumption were observed in the growing period (29 to 55 kg liveweight), the finishing period (55 to 90 kg liveweight), the 'transition' period (90 to 106 kg liveweight, group-feeding), or the pre-gestation period (106 to 116 kg liveweight, individually fed to the time of breeding). However, pigs fed rapeseed meal consistently consumed slightly less feed during the growing, finishing and transition periods. Manns (1962) and Hussar (1958) observed a depression in feed intake of group-fed pigs receiving rapeseed meal.

At the beginning of the pre-gestation period, lot 3 was changed from the rapeseed meal diet to the soybean meal diet, and lot 4 was changed from the soybean meal diet to the rapeseed meal diet. All lots were individually fed at this time, and feed intake was restricted to 2.7 kg daily. On individual feeding, few pigs consumed their allotted feed and average daily feed was 1.8 kg. In the first 2 weeks on this feeding schedule, feed intake of lot 3 pigs was lower than for all other groups. For lot 4 (changed from the soybean meal to the rapeseed meal diet), feed intake during this same period was higher than for all other

THE JOURNAL OF THE

THE JOURNAL OF THE

THE JOURNAL OF THE

THE JOURNAL OF THE

THE JOURNAL OF THE

THE JOURNAL OF THE

groups. This indicated that the palatability of rapeseed meal was not a problem, but that conversely, changing from rapeseed meal diets to soybean meal diets may give problems in acceptability. Feed consumption during gestation will be discussed under reproductive performance.

Table 6. Average daily feed, average daily gain, and efficiency of feed utilization -- experiment 1.

Lot	Feeding period	Rapeseed meal for ¹		No. of pigs	Avg daily feed kg	Avg daily gain gm	Feed/kg gain
		growth	reproduction				
1	Growing	-	-	6	2.08	695	2.99
2		+	+	6	2.04	663	3.08
3		+	-	6	2.00	705	2.84
4		-	+	6	2.08	709	2.94
1	Finishing	-	-	6	3.17	914	3.48
2		+	+	6	3.42	895	3.84
3		+	-	6	2.96	809	3.68
4		-	+	6	3.73	954	3.91
1	Growing and Finishing	-	-	6	2.68	817	3.29
2		+	+	6	2.81	791	3.56
3		+	-	6	2.55	763	3.35
4		-	+	6	2.97	840	3.54
1	Transition-group-fed	-	-	6	2.82	413	6.82
2		+	+	6	2.88	350	8.23
3		+	-	6	2.67	322	8.23
4		-	+	6	3.01	454	6.60
1	Pre-gestation -individually fed	-	-	6	1.86	160	- ²
2		+	+	6	1.73	128	-
3		+	-	6	1.71	172	-
4		-	+	6	1.89	188	-

¹+ indicates a level of 8% rapeseed meal in the diet and - indicates no rapeseed meal

²- these values were not determined.

Table 7. Treatment means for average daily feed, average daily gain and efficiency of feed utilization -- experiment 1

Feeding period	Treatment Diet to 89 kg	No. of pigs	Avg daily feed kg	Avg daily gain gm	Feed/ kg gain kg
Growing	SM	12	2.08	702	2.96
	RM	12	2.02	684	2.96
Finishing	SM	12	3.45	934	3.70
	RM	12	3.19	852 ⁺	3.76
Growing and finishing	SM	12	2.78	828	3.48
	RM	12	2.68	777 ⁺	3.46
Transition ¹	SM	12	2.92	434	6.71
	RM	12	2.78	336*	8.23*
Pre-gestation	SM	12	1.88	174	-
	RM	12	1.72	150	-

¹Pigs were group-fed in the transition period and individually fed in the pre-gestation period. At the end of this period, one lot was changed from the basal to the rapeseed meal diet, one lot was changed from the rapeseed meal diet to the basal diet and two lots were not changed.

⁺P < 0.10

^{*}P < 0.05

b) Average daily gain

In the growing period, gilts fed rapeseed meal grew 18 gm per day less than those fed soybean meal (not significantly different). In the finishing period, pigs fed rapeseed meal grew 82 gm per day less than those fed soybean meal ($P < 0.10$), and this appeared to be associated with a decreased feed intake. In the combined growing and finishing periods, the gilts grew 51 gm daily less ($P < 0.10$) when fed 8% rapeseed meal as compared to those fed the basal soybean meal diets. This

1. The following are the results of the analysis of the data obtained from the experiments conducted during the period from 1950 to 1952.

Year	1950	1951	1952	Total	Percentage of Total	Remarks
1950	100	100	100	300	100%	
1951	100	100	100	300	100%	
1952	100	100	100	300	100%	
Total	300	300	300	900	100%	
1950	100	100	100	300	100%	
1951	100	100	100	300	100%	
1952	100	100	100	300	100%	
Total	300	300	300	900	100%	
1950	100	100	100	300	100%	
1951	100	100	100	300	100%	
1952	100	100	100	300	100%	
Total	300	300	300	900	100%	

The following are the results of the analysis of the data obtained from the experiments conducted during the period from 1950 to 1952.

1950 100
1951 100
1952 100

CONCLUSIONS

The following are the results of the analysis of the data obtained from the experiments conducted during the period from 1950 to 1952.

depression of growth is in agreement with the findings of Manns (1962).

In the transition period (from 90 to 106 kg liveweight), average daily gain of gilts fed rapeseed meal was 98 gm less ($P < 0.05$) than of those fed soybean meal diets. The reason for this difference was not evident, but possibly the stress of moving and changing environment affected more severely pigs fed rapeseed meal. When the feed was changed and individual feeding was begun (pre-gestation period), those changed to the soybean meal diet from the rapeseed meal diet did not gain or lose weight on the average in the first 2 weeks on this diet (4 out of 6 pigs lost weight), while those changed to rapeseed meal diets from the soybean meal diets gained on the average 180 gm daily in the same period. However, during the whole pre-gestation period (106 kg to breeding time), sows fed rapeseed meal for growth and rapeseed meal for reproduction (R-R*) grew most slowly.

c) Efficiency of feed utilization

Efficiency of feed utilization, measured as kg of feed per kg of gain was similar for pigs fed rapeseed meal and those fed soybean meal for the growing and finishing periods. During the growing period all pigs consumed 2.96 kg of feed per kg of gain and during the finishing period, they consumed 3.73 kg of feed per kg of gain. In the transition period, pigs fed rapeseed meal required an additional 1.52 kg of feed per kg gain than was required by pigs fed soybean meal, a difference of 23% more feed per kg gain ($P < 0.05$). During pre-gestation, feed efficiency was not calculated since several^{*} animals lost weight and values

*The lots are designated as B-B, R-R, R-B, and B-R as follows: B-B, soybean meal diet (basal) for growth and for reproduction; R-R, rapeseed meal diet for growth and for reproduction; R-B, rapeseed meal diet for growth and soybean meal diet for reproduction; and B-R, soybean meal diet for growth and rapeseed meal diet for reproduction.

obtained would have little meaning.

2. Reproductive performance

Tables 8, 9 and 10 list the means of the reproductive and breeding traits measured for the first litter, for the second litter and treatment means for diet and litter, respectively.

a) First litter

When the sows were bred for the first time, two sows failed to conceive. These sows were littermates; one was in lot 2 (R-R) and one was in lot 3 (R-B). The sows were slaughtered, and examination of the reproductive tracts* revealed that no corpora lutea were present, and ovaries appeared to be smaller than normal. It was concluded that the reproductive tracts were infantile; that is, not fully developed.

During gestation, daily feed was restricted to 2.3 kg, and consumption was 2.2 kg daily on the average, with little variation between groups. Average daily gain was also similar for all groups. No differences between lots in feed efficiency which averaged 4.54 kg feed per kg gain, were observed.

The breeding performance of sows in this study was not affected by the feeding of rapeseed meal. The number of services to conception, weight at breeding, gestation gains, lactation losses, and the change in weight of the sows from breeding to weaning of the litters were not affected by dietary treatments. These results differ from those of Manns (1962) who showed that sows fed rapeseed meal had to be bred 2 to 2.5 times before conception occurred.

At farrowing time, one sow in lot 1(B-B) farrowed 5 dead pigs and died 2 days later due to unknown causes. Post-mortem findings showed

*The advice of Dr. T.D.D. Groves, Assistant Professor of Physiology, is acknowledged.

Washington, D.C. 20540

January 10, 1964

Dear Mr. [Name]:
I am pleased to hear from you and to learn that you are
interested in the [Project Name] project. I am sure that
you will find the information I am about to provide to you
very helpful.

Very truly yours,

[Name]
[Title]
[Organization]
[Address]
[City]
[State]
[Zip]

cc: [Name]

[Name]
[Title]
[Organization]
[Address]
[City]
[State]
[Zip]

2 macerated feti in the uterus. A sow in lot 4 (B-R), farrowing on the same date (June 24), farrowed 3 dead pigs, and died 3 days later. A tear in the uterine wall was found on post-mortem examination, and 6 piglets were found in the abdominal cavity. The cause of death was severe peritonitis. The loss of 4 sows, 2 due to non-conception and 2 due to death of the sows at farrowing time, left the lots balanced with 5 sows per lot.

Table 8. Reproductive performance -- first litters, experiment 1

Lot		1(B-B)	2(R-R)	3(R-B)	4(B-R)
Growing diet		SM	RM	RM	SM
Reproduction diet		SM	RM	SM	RM
Reproductive traits					
Number of litters ¹		5	5	5	5
<u>Gestation data</u>					
Number of services to conception		1.0	1.0	1.4	1.2
Breeding weight	kg	116	114	117	120
Average daily feed	kg	2.18	2.10	2.21	2.23
Average daily gain	kg	0.47	0.50	0.49	0.49
Feed per kg gain	kg	4.56	4.28	4.49	4.64
Total gestation gain	kg	56.1	52.4	51.9	48.6
Length of gestation	days	115	115	115	114
<u>Lactation data</u>					
Feed during lactation	kg	79.4	67.6	65.5	84.0
Lactation loss	kg	13.5	19.3	11.0	12.0
Increase in sow's weight (breeding to weaning of litters)	kg	26.5	22.2	26.1	22.7
<u>Litter data</u>					
Number of pigs born		10.2	10.6	9.0	9.8
Number of live pigs born		9.8 ^{a,b}	10.0 ^a	7.8 ^b	9.4 ^{a,b}
Number of pigs weaned		8.0 ^{a,b}	6.2 ^{a,b}	5.6 ^a	8.8 ^b
Litter birth weight	kg	13.5	12.2	11.5	12.2
Litter weaning weight	kg	41.1	26.7	31.0	41.3
Avg pig birth weight	kg	1.36	1.23	1.48	1.42
Avg pig weaning weight	kg	5.13	4.25	5.60	5.06

¹One sow failed to farrow in each lot. Sows in lots 2 and 3 that failed to farrow did not conceive.

^{a,b}Figures with different superscripts are different at $P < 0.05$ as determined by Duncan's multiple range test (Steele and Torrie, 1960).

Table 9. Reproductive performance -- second litters, experiment 2

Lot		1(B-B)	2(R-R)	3(R-B)	4(B-R)
Growing diet		SM	RM	RM	SM
Reproduction diet		SM	RM	SM	RM
<u>Reproductive traits¹</u>					
Number of litters ²		4	4	4	4
<u>Gestation data</u>					
Number of services to conception		1.25	2.0	1.0	1.0
Breeding weight	kg	145	148	144	140
Total gestation gain	kg	46.8	48.8	41.0	46.0
Length of gestation	days	115	114	115	115
<u>Lactation data</u>					
Lactation loss	kg	15.3	18.6	14.7	15.3
Increase in sows' weight (breeding to weaning of litters)	kg	22.7	22.1	21.2	24.3
<u>Litter data</u>					
Number of pigs born		10.8	10.0	9.5	10.0
Number of live pigs born		10.0	9.8	9.5	9.5
Number of pigs weaned		9.5	7.8	8.8	8.8
Litter birth weight	kg	14.8	14.1	14.9	13.6
Litter weaning weight	kg	50.0	46.5	51.7	45.8
Avg pig birth weight	kg	1.46	1.45	1.58	1.45
Avg pig weaning weight	kg	5.28	5.25	5.94	5.34

¹No significant differences were observed for any traits.

²Results for one litter in lot 1 and one in lot 4 were removed so that the number of litters per lot was balanced. One sow in lot 2 and one in lot 3 failed to conceive.

THE UNIVERSITY OF CHICAGO

1910

1910

1910

1910

1910

1910

1910

1910

1910

1910

1910

1910

1910

1910

1910

1910

1910

1910

1910

1910

1910

1910

1910

1910

1910

1910

1910

1910

1910

1910

1910

1910

1910

1910

Table 10. Treatment means for reproductive performance

Reproductive traits	Litter no.		Growing diet		Reproduction diet		Mean
	1	2	SM	RM	SM	RM	
Number of litters ¹	16	16	16	16	16	16	32
<u>Gestation data</u>							
Number of services to conception	1.1	1.3	1.1	1.2	1.1	1.2	1.2
Breeding weight	kg 117	144	130	131	130	131	130
Gestation gain	kg 53.0	45.6	49.7	49.0	49.3	49.5	49.3
<u>Lactation data</u>							
Lactation loss	kg 14.9	16.0	14.5	16.4	12.8	18.1	15.4
Increase in sows' weight (breeding to weaning of litters)	kg 24.9	22.6	24.8	22.7	25.4	22.2	+23.7
<u>Litter data</u>							
Number of pigs born	10.2	10.1	10.2	10.0	9.8	10.5	10.1
Number of live pigs born	9.3	9.7	9.6	9.4	9.2	9.8	9.5
Number of pigs weaned	7.2*	8.7	8.9	7.0	8.0	8.0	8.0
Litter birth weight	kg 12.2 ⁺	14.3	13.3	13.2	13.4	13.2	13.2
Litter weaning weight	kg 34.7**	48.5	44.5*	38.6	43.2	40.0	41.6
Avg pig birth weight	kg 1.32*	1.48	1.40	1.42	1.46	1.35	1.40
Avg pig weaning weight	kg 4.92*	5.45	5.16	5.21	5.48	4.90	5.19

⁺ P < 0.10

* P < 0.05

**P < 0.01

¹Data are included only for sows farrowing in both the first and second reproductive cycles.

A difference between 2 lots (P < 0.05) in the number of live pigs farrowed was observed. The R-R lot (lot 2) farrowed 2.2 live pigs less per litter than the R-B lot (lot 3).

A difference (P < 0.05) associated with dietary treatments during growth was found in the number of pigs weaned when these values were adjusted by analysis of covariance for the number of live pigs born. The sows fed rapeseed meal diets during growth (lots 2 and 3) weaned, on the average 5.9 pigs per litter and those fed soybean meal diets for this

period weaned 8.4 pigs per litter. These results suggest that the feeding of rapeseed meal during growth impaired future milk producing ability of sows. This is in general agreement with the observations of Manns (1962) who suggested that lactation was impaired by the feeding of rapeseed meal to females during growth and reproduction. For the first litter, a difference in litter weaning weights ($P < 0.05$) was observed. This difference amounted to 12.4 kg per litter or 30% lower litter weaning weights for the sows fed rapeseed meal during growth.

No differences were observed in total number of pigs born, litter birth weight, average pig weaning weight and average pig birth weight associated with the feeding of 8% rapeseed meal during growth or reproduction or both. Bell and Belzile (1965) reported that dietary rapeseed meal led to a depression in number of pigs born alive, birth weight and weaning weight of litters. Manns and Bowland (1963a) observed that the number of pigs born per litter was similar for sows fed rapeseed meal and those fed soybean meal diets.

b) Second litter

When rebred, the sows conceived an average of 10 days after weaning of the first litters. One sow in lot 2 (R-R) and one in lot 3 (R-B) failed to conceive when bred for 3 consecutive estrous cycles. These sows were in the same lots in which sows failed to conceive in the first reproductive cycle, and in this case again the two sows were littermates. The reproductive tracts of these two sows were examined. In the sow from lot 2, apparently no follicles had ruptured in the previous 5 weeks, and no corpora lutea were observed on either ovary. The ovaries and uterus of both sows appeared normal. The failure of 4 sows to conceive in two reproductive cycles in lots fed rapeseed meal for growth suggests

impairment of reproductive ability associated with feeding of rapeseed meal to young female swine.

Reproductive performance for the second litters was analyzed by removing data for one sow in lot 1 and one sow in lot 4, both of which were littermates of the sows failing to conceive, leaving the groups balanced with 4 litters per lot for statistical analysis. When the lots were balanced this way, no differences in measured reproductive traits were observed in the second litter. Any harmful effects of rapeseed meal appear to affect gilts more than sows in their second litter and may, as Kennedy and Purves (1941) suggested for rats, be associated with a delay in ovary development. These data also suggest that certain litter groups may be affected more than others.

c) First and second litters

Average birth weight, average weaning weight and numbers of pigs weaned were greater ($P < 0.05$) for the second than for the first litter, and litter weaning weights were higher ($P < 0.01$). Gestation gain for the first litter was greater than for the second ($P < 0.05$). No differences between first and second litters were observed in number of live pigs farrowed, lactation losses of the sows, and changes in sows' weights from breeding to weaning of litters. Birth weight of the second litter was greater at the 10% level of significance.

3. Lactation studies

Tables 11, 12, 13 and 14 list, respectively, the gross composition of sows' colostrum and milk, treatment means for gross colostrum and milk composition, fatty acid composition of sows' colostral and milk fat and treatment means for fatty acid composition.

Table 11. Gross composition of colostrum and milk

Lot		1	2	3	4
Feed for growth		SM	RM	RM	SM
Feed for reproduction		SM	RM	SM	RM
<u>Colostrum</u>					
Total solids	%	20.67 ^b	23.33 ^{a,b}	24.12 ^a	23.61 ^{a,b}
Fat	%	6.40 ^b	7.71 ^{a,b}	8.37 ^a	6.72 ^{a,b}
Solids-not-fat	%	14.27	15.62	15.88	16.94
Protein	%	8.80 ^b	10.85 ^{a,b}	10.43 ^{a,b}	11.59 ^a
Lactose	%	4.81	4.14	4.65	4.69
Ash	%	0.66	0.63	0.67	0.61
<u>Milk</u>					
Total solids	%	18.49	20.82	19.50	18.59
Fat	%	6.83	8.06	7.61	6.17
Solids-not-fat	%	11.66	12.77	11.83	12.41
Protein	%	5.25	5.48	5.63	5.35
Lactose	%	5.70	6.49	5.49	6.33
Ash	%	0.71	0.79	0.77	0.74

^{ab}Numbers with different superscripts are different at $P < 0.05$.

Table 1. Summary of the experimental results (continued)

Run	Time (min)	Temp (°C)	Pressure (mm Hg)	Flow rate (ml/min)	Yield (%)
1	10	100	1.0	1.0	100
2	20	100	1.0	1.0	100
3	30	100	1.0	1.0	100
4	40	100	1.0	1.0	100
5	50	100	1.0	1.0	100
6	60	100	1.0	1.0	100
7	70	100	1.0	1.0	100
8	80	100	1.0	1.0	100
9	90	100	1.0	1.0	100
10	100	100	1.0	1.0	100
11	110	100	1.0	1.0	100
12	120	100	1.0	1.0	100
13	130	100	1.0	1.0	100
14	140	100	1.0	1.0	100
15	150	100	1.0	1.0	100
16	160	100	1.0	1.0	100
17	170	100	1.0	1.0	100
18	180	100	1.0	1.0	100
19	190	100	1.0	1.0	100
20	200	100	1.0	1.0	100
21	210	100	1.0	1.0	100
22	220	100	1.0	1.0	100
23	230	100	1.0	1.0	100
24	240	100	1.0	1.0	100
25	250	100	1.0	1.0	100
26	260	100	1.0	1.0	100
27	270	100	1.0	1.0	100
28	280	100	1.0	1.0	100
29	290	100	1.0	1.0	100
30	300	100	1.0	1.0	100
31	310	100	1.0	1.0	100
32	320	100	1.0	1.0	100
33	330	100	1.0	1.0	100
34	340	100	1.0	1.0	100
35	350	100	1.0	1.0	100
36	360	100	1.0	1.0	100
37	370	100	1.0	1.0	100
38	380	100	1.0	1.0	100
39	390	100	1.0	1.0	100
40	400	100	1.0	1.0	100
41	410	100	1.0	1.0	100
42	420	100	1.0	1.0	100
43	430	100	1.0	1.0	100
44	440	100	1.0	1.0	100
45	450	100	1.0	1.0	100
46	460	100	1.0	1.0	100
47	470	100	1.0	1.0	100
48	480	100	1.0	1.0	100
49	490	100	1.0	1.0	100
50	500	100	1.0	1.0	100

Notes: (a) The reaction was carried out in a 100 ml round-bottomed flask equipped with a magnetic stirrer and a nitrogen inlet.

Table 12. Treatment means for gross composition of colostrum and milk

		Growing diet		Reproduction diet		Colostrum	Milk	Mean
		SM	RM	SM	RM			
<u>Colostrum</u>								
Total solids	%	22.14	23.72 ⁺	22.40	23.47	22.93*	19.35	21.18
Fat	%	6.56	8.04*	7.38	7.22	7.30	7.17	7.24
Solids-not-fat	%	15.60	15.75	15.08	16.33	15.68**	12.17	13.92
Protein	%	10.20	10.64	9.62	11.22	10.42**	5.43	7.92
Lactose	%	4.75	4.40	4.73	4.42	4.57**	6.00	5.28
Ash	%	0.63	0.65	0.66	0.62	0.64**	0.75	0.70
<u>Milk</u>								
Total solids	%	18.54	20.16	19.00	19.70			
Fat	%	6.50	7.84	7.22	7.11			
Solids-not-fat	%	12.04	12.30	11.74	12.59			
Protein	%	5.30	5.56	5.54	5.42			
Lactose	%	6.02	5.99	5.60	6.41			
Ash	%	0.73	0.78	0.74	0.76			

⁺P < 0.10

*P < 0.05

**P < 0.01

TABLE I. Summary of the results of the calculations for the various cases considered. The values in parentheses are the values of the parameters used in the calculations.

Case		Parameters		Results		Comments
No.	Label	α	β	γ	δ	
1	(a)	0.1	0.1	0.1	0.1	Good
2	(b)	0.2	0.2	0.2	0.2	Good
3	(c)	0.3	0.3	0.3	0.3	Good
4	(d)	0.4	0.4	0.4	0.4	Good
5	(e)	0.5	0.5	0.5	0.5	Good
6	(f)	0.6	0.6	0.6	0.6	Good
7	(g)	0.7	0.7	0.7	0.7	Good
8	(h)	0.8	0.8	0.8	0.8	Good
9	(i)	0.9	0.9	0.9	0.9	Good
10	(j)	1.0	1.0	1.0	1.0	Good
11	(k)	0.1	0.2	0.3	0.4	Good
12	(l)	0.2	0.3	0.4	0.5	Good
13	(m)	0.3	0.4	0.5	0.6	Good
14	(n)	0.4	0.5	0.6	0.7	Good
15	(o)	0.5	0.6	0.7	0.8	Good
16	(p)	0.6	0.7	0.8	0.9	Good
17	(q)	0.7	0.8	0.9	1.0	Good
18	(r)	0.8	0.9	1.0	1.1	Good
19	(s)	0.9	1.0	1.1	1.2	Good
20	(t)	1.0	1.1	1.2	1.3	Good

TABLE I
continued

Table 13. Fatty acid composition of sows' colostrum and milk fat

Lot	Colostrum fat				Milk fat			
	1	2	3	4	1	2	3	4
Feed for growth	SM	RM	RM	SM	SM	RM	RM	SM
Feed for reproduction	SM	RM	SM	RM	SM	RM	SM	RM
<u>Fatty acids-% of total</u>								
C 10:0 (capric)	- ¹	-	-	-	0.16	0.15	0.18	0.26
C 12:0 (lauric)	-	-	-	-	0.22	0.23	0.24	0.34
C 14:0 (myristic)	1.52	1.40	1.58	1.62	2.92 ^b	3.07 ^b	3.46 ^{ab}	3.87 ^a
C 14:1 (myristoleic)	0.04	0.06	0.11	0.08	0.11	0.14	0.15	0.23
C 16:0 (palmitic)	22.51	21.70	22.71	23.05	26.70	26.52	26.64	29.34
C 16:1 (palmitoleic)	5.18	5.16	5.10	4.63	7.74 ^b	7.58 ^b	8.89 ^{ab}	10.29 ^a
C 18:0 (stearic)	5.16	4.52	4.99	5.51	4.72	4.34	4.09	4.06
C 18:1 (oleic)	41.14	44.05	39.91	39.68	41.39	43.57	40.53	36.67
C 18:2 (linoleic)	19.77	19.72	20.46	21.17	13.26	11.57	12.45	11.81
C 20:0 ² (arachidic)	1.81	1.58	1.85	1.51	0.99	1.04	1.10	1.05
C 20:4 (arachidonic)	0.94	-	0.90	0.28	0.68	0.26	0.76	0.58
C 14:0, 16:0 & 18:0	30.64	28.59	30.91	31.55	35.47	35.10	35.60	39.84
C 16:1, 18:1 & 18:2	69.36	71.41	69.09	68.45	64.53	64.90	64.40	60.16

^{ab}Values with different superscripts differ at $P < 0.05$.

¹Levels were not detectable.

²C 18:3 peak covered by this peak on chromatogram, so this value includes both C 18:3 and C 20:0 fatty acids.

THE JOURNAL OF THE AMERICAN MEDICAL ASSOCIATION PUBLISHED WEEKLY

CONTENTS				ORIGINAL ARTICLES				DEPARTMENTS	
1	2	3	4	5	6	7	8	9	10
11	12	13	14	15	16	17	18	19	20
21	22	23	24	25	26	27	28	29	30
31	32	33	34	35	36	37	38	39	40
41	42	43	44	45	46	47	48	49	50
51	52	53	54	55	56	57	58	59	60
61	62	63	64	65	66	67	68	69	70
71	72	73	74	75	76	77	78	79	80
81	82	83	84	85	86	87	88	89	90
91	92	93	94	95	96	97	98	99	100
101	102	103	104	105	106	107	108	109	110
111	112	113	114	115	116	117	118	119	120
121	122	123	124	125	126	127	128	129	130
131	132	133	134	135	136	137	138	139	140
141	142	143	144	145	146	147	148	149	150
151	152	153	154	155	156	157	158	159	160
161	162	163	164	165	166	167	168	169	170
171	172	173	174	175	176	177	178	179	180
181	182	183	184	185	186	187	188	189	190
191	192	193	194	195	196	197	198	199	200
201	202	203	204	205	206	207	208	209	210
211	212	213	214	215	216	217	218	219	220
221	222	223	224	225	226	227	228	229	230
231	232	233	234	235	236	237	238	239	240
241	242	243	244	245	246	247	248	249	250
251	252	253	254	255	256	257	258	259	260
261	262	263	264	265	266	267	268	269	270
271	272	273	274	275	276	277	278	279	280
281	282	283	284	285	286	287	288	289	290
291	292	293	294	295	296	297	298	299	300
301	302	303	304	305	306	307	308	309	310
311	312	313	314	315	316	317	318	319	320
321	322	323	324	325	326	327	328	329	330
331	332	333	334	335	336	337	338	339	340
341	342	343	344	345	346	347	348	349	350
351	352	353	354	355	356	357	358	359	360
361	362	363	364	365	366	367	368	369	370
371	372	373	374	375	376	377	378	379	380
381	382	383	384	385	386	387	388	389	390
391	392	393	394	395	396	397	398	399	400
401	402	403	404	405	406	407	408	409	410
411	412	413	414	415	416	417	418	419	420
421	422	423	424	425	426	427	428	429	430
431	432	433	434	435	436	437	438	439	440
441	442	443	444	445	446	447	448	449	450
451	452	453	454	455	456	457	458	459	460
461	462	463	464	465	466	467	468	469	470
471	472	473	474	475	476	477	478	479	480
481	482	483	484	485	486	487	488	489	490
491	492	493	494	495	496	497	498	499	500
501	502	503	504	505	506	507	508	509	510
511	512	513	514	515	516	517	518	519	520
521	522	523	524	525	526	527	528	529	530
531	532	533	534	535	536	537	538	539	540
541	542	543	544	545	546	547	548	549	550
551	552	553	554	555	556	557	558	559	560
561	562	563	564	565	566	567	568	569	570
571	572	573	574	575	576	577	578	579	580
581	582	583	584	585	586	587	588	589	590
591	592	593	594	595	596	597	598	599	600
601	602	603	604	605	606	607	608	609	610
611	612	613	614	615	616	617	618	619	620
621	622	623	624	625	626	627	628	629	630
631	632	633	634	635	636	637	638	639	640
641	642	643	644	645	646	647	648	649	650
651	652	653	654	655	656	657	658	659	660
661	662	663	664	665	666	667	668	669	670
671	672	673	674	675	676	677	678	679	680
681	682	683	684	685	686	687	688	689	690
691	692	693	694	695	696	697	698	699	700
701	702	703	704	705	706	707	708	709	710
711	712	713	714	715	716	717	718	719	720
721	722	723	724	725	726	727	728	729	730
731	732	733	734	735	736	737	738	739	740
741	742	743	744	745	746	747	748	749	750
751	752	753	754	755	756	757	758	759	760
761	762	763	764	765	766	767	768	769	770
771	772	773	774	775	776	777	778	779	780
781	782	783	784	785	786	787	788	789	790
791	792	793	794	795	796	797	798	799	800
801	802	803	804	805	806	807	808	809	810
811	812	813	814	815	816	817	818	819	820
821	822	823	824	825	826	827	828	829	830
831	832	833	834	835	836	837	838	839	840
841	842	843	844	845	846	847	848	849	850
851	852	853	854	855	856	857	858	859	860
861	862	863	864	865	866	867	868	869	870
871	872	873	874	875	876	877	878	879	880
881	882	883	884	885	886	887	888	889	890
891	892	893	894	895	896	897	898	899	900
901	902	903	904	905	906	907	908	909	910
911	912	913	914	915	916	917	918	919	920
921	922	923	924	925	926	927	928	929	930
931	932	933	934	935	936	937	938	939	940
941	942	943	944	945	946	947	948	949	950
951	952	953	954	955	956	957	958	959	960
961	962	963	964	965	966	967	968	969	970
971	972	973	974	975	976	977	978	979	980
981	982	983	984	985	986	987	988	989	990
991	992	993	994	995	996	997	998	999	1000

Published by the American Medical Association, 535 North Dearborn Street, Chicago, Ill. 60610.
 Second-class postage paid at Chicago, Ill., and at additional mailing offices.
 Postmaster: This publication is entered as second-class matter, October 3, 1917, under post office number 384, at Chicago, Ill., and at additional mailing offices. Postage paid at Chicago, Ill., and at additional mailing offices.
 Copyright, 1961, by American Medical Association.
 Printed at the American Medical Association Press, Chicago, Ill.

Table 14. Treatment means for fatty acid composition of colostrum and milk fat

Fatty acids % of total	Growing diet		Reproduction diet		Colostrum**	Milk	Mean
	SM	RM	SM	RM			
C 10:0 (milk fat only)	0.21	0.16	0.17	0.20	0.00 ¹	0.19	
C 12:0 (milk fat only)	0.28	0.24	0.23	0.28	0.00	0.26	
C 14:0	2.48	2.38	2.38	2.48	1.53	3.33	2.43
C 14:1	0.11	0.12	0.10	0.13	0.07	0.16	0.12
C 16:0	25.40	24.39	24.64	25.15	22.49	27.30	24.90
C 16:1	6.95	6.68	6.73	6.91	5.02	8.62	6.82
C 18:0	4.86	4.48	4.74	4.60	5.04	4.30	4.67
C 18:1	39.72	42.01	40.74	41.00	41.19	40.54	40.87
C 18:2	16.50	16.04	16.49	16.06	20.28	12.27	16.28
C 20:0 ²	1.34	1.39	1.44	1.30	1.69	1.04	1.37
C 20:4	0.62	0.48	0.82	0.28	0.53	0.57	0.55
C 14:0, 16:0 & 18:0	34.38	32.54	33.16	33.77	30.42	36.50	33.46
C 16:1, 18:1 & 18:2	65.62	67.46	66.84	66.23	69.58	63.50	66.54

**P < 0.01

¹Levels were not detectable.

²C 18:3 peak covered by this peak on chromatogram, so this value includes both C 18:3 and C 20:0 fatty acids.

a) Gross colostrum and milk composition

For the colostrum, a difference of 3.4% in total solids ($P < 0.05$) was observed between the B-B group (20.7%) and the R-B group (24.1%). However, the R-R group was not different from the B-B group, so the differences observed in total solids cannot be attributed directly to diet. Percent fat in the B-B group was lower ($P < 0.05$) at 6.4% than that of the R-B group, at 8.4%. In the two lots fed rapeseed meal for growth (lots 2 and 3), percent fat was higher than for those fed soybean meal diets (8.0 compared to 6.6%). No differences associated with diet were observed in percent solids-not-fat (SNF), protein, ash or lactose of sows' colostrum.

Compared to colostrum, sows' milk had lower levels of total solids ($P < 0.05$), SNF, and protein and higher levels of ash ($P < 0.01$ for the last three components). No differences in percent fat were observed between milk and colostrum of the sows. In sows' milk, no significant differences associated with diet were observed between lots in any of the components measured. Percent fat and percent solids appeared to be higher in milk of sows fed rapeseed meal for growth.

b) Fatty acid composition

No differences due to dietary treatments were observed in fatty acid levels in the fat of colostrum. Percent saturated fat and percent unsaturated fat did not vary between lots. Only traces of C 10:0 (capric) and C 12:0 (lauric) fatty acids were observed in the colostrum fat of sows, which agrees with the results of DeMan and Bowland (1963) and Stinson (1966).

All fatty acid levels measured in sows' milk fat were different from those of colostrum ($P < 0.01$), with the exception of C 18:1

THE HISTORY OF THE UNITED STATES

The first of these is the fact that the United States is a young nation, and that its history is a history of growth and expansion. The second is the fact that the United States is a nation of immigrants, and that its history is a history of the struggle for the rights of these immigrants. The third is the fact that the United States is a nation of free men, and that its history is a history of the struggle for the rights of these free men. The fourth is the fact that the United States is a nation of law, and that its history is a history of the struggle for the rights of these laws. The fifth is the fact that the United States is a nation of peace, and that its history is a history of the struggle for the rights of these peace. The sixth is the fact that the United States is a nation of justice, and that its history is a history of the struggle for the rights of these justice. The seventh is the fact that the United States is a nation of liberty, and that its history is a history of the struggle for the rights of these liberty. The eighth is the fact that the United States is a nation of equality, and that its history is a history of the struggle for the rights of these equality. The ninth is the fact that the United States is a nation of unity, and that its history is a history of the struggle for the rights of these unity. The tenth is the fact that the United States is a nation of progress, and that its history is a history of the struggle for the rights of these progress.

THE HISTORY OF THE UNITED STATES

The first of these is the fact that the United States is a young nation, and that its history is a history of growth and expansion. The second is the fact that the United States is a nation of immigrants, and that its history is a history of the struggle for the rights of these immigrants. The third is the fact that the United States is a nation of free men, and that its history is a history of the struggle for the rights of these free men. The fourth is the fact that the United States is a nation of law, and that its history is a history of the struggle for the rights of these laws. The fifth is the fact that the United States is a nation of peace, and that its history is a history of the struggle for the rights of these peace. The sixth is the fact that the United States is a nation of justice, and that its history is a history of the struggle for the rights of these justice. The seventh is the fact that the United States is a nation of liberty, and that its history is a history of the struggle for the rights of these liberty. The eighth is the fact that the United States is a nation of equality, and that its history is a history of the struggle for the rights of these equality. The ninth is the fact that the United States is a nation of unity, and that its history is a history of the struggle for the rights of these unity. The tenth is the fact that the United States is a nation of progress, and that its history is a history of the struggle for the rights of these progress.

(oleic) - the major component, and C 20:4. Diet did not have a significant effect on fatty acid levels in the fat of sows' milk. C 14:1 and C 14:0 appeared to be higher in fat of the milk of lot 4 (B-R) sows than that of lots 2 and 3 sows, which received rapeseed meal diets during the growth periods. Levels of saturated fatty acids appeared to be higher (not significant) in the milk fat of lot 4 sows than that of other groups. Saturated fatty acids were 39% of the fat for lot 4 and 35% of milk fat for the other 3 lots.

B. Progeny Growth Studies -- Experiment 2

1. Feed consumption, daily gain and efficiency of feed utilization

Table 15 shows the lot means for feed consumption, average daily gain and efficiency of feed utilization for experiment 2. Treatment means are listed in table 16.

a) Average daily feed consumption

Dietary rapeseed meal led to a depression ($P \leq 0.05$) of 280 gm feed per day in the finishing period (55 kg to 89 kg liveweight) as compared to dietary soybean meal. In the starting period (6 kg to 18 kg liveweight), pigs fed rapeseed meal ate 4.7% less than those fed soybean meal diets, in the growing period (18 to 55 kg liveweight), 1.6% less, and in the combined starting, growing and finishing periods, 3.4% less, none of which were significantly different. These observations are in agreement with earlier observations in this report on growing gilts. Hussar and Bowland (1959a) and Seale in 1952 (cited by Bowland, 1965a), reported variable effects of rapeseed meal on feed consumption, although Manns and Bowland (1963a) reported that rapeseed meal did not affect feed intake.

Male pigs consumed significantly more feed than females ($P \leq 0.05$). The multiple range test showed that males fed rapeseed meal diets gained

less than males fed soybean meal diets in the finishing period ($P < 0.05$). However, significant interactions were not observed between sex and dietary treatment when compared by an F-test in analysis of variance.

b) Average daily gain

Pigs fed soybean meal diets grew 10% more rapidly throughout the period from 6 to 89 kg liveweight as shown by a difference of 64 gm average daily gain when compared to pigs fed rapeseed meal diets ($P < 0.05$). This observation is in agreement with observations of Manns and Bowland (1963a), Hussar and Bowland (1959a) and Seale in 1952 (cited by Bowland, 1965a), who noted a depression in average daily gain in swine on levels of rapeseed meal from 8% to 15% of the diet. This depression is consistent with observations earlier in this report that young female pigs from 29 kg to 89 kg liveweight grew at a rate of 50 gm less daily when fed rapeseed meal in substitution for soybean meal in their diets. Compared to pigs fed soybean meal, in the starting period average daily gain of pigs fed rapeseed meal was depressed 14% (58 gm daily), in the growing period gain was depressed 9% (66 gm daily, $P < 0.10$), and in the finishing period gain was depressed 9% (81 gm daily, $P < 0.05$).

The offspring of sows fed rapeseed meal gained as rapidly as the offspring of sows fed soybean meal as the main protein source, throughout the period from 6 to 89 kg liveweight. An average daily gain of 674 gm was observed for progeny of sows fed soybean meal as compared to 658 gm daily for progeny of sows fed rapeseed meal, a difference of 16 gm. However, in the starting period (6 to 18 kg liveweight), gain was 40 gm less daily for offspring of R-R sows than for offspring of B-B sows and in the growing period (18 to 55 kg liveweight), 45 gm less. In the finishing period (55 to 89 kg liveweight), 31 gm more daily was

The first of these is the fact that the system is not in equilibrium. The second is the fact that the system is not in equilibrium. The third is the fact that the system is not in equilibrium.

THE SECOND OF THESE

The second of these is the fact that the system is not in equilibrium. The third is the fact that the system is not in equilibrium. The fourth is the fact that the system is not in equilibrium.

The third of these is the fact that the system is not in equilibrium. The fourth is the fact that the system is not in equilibrium. The fifth is the fact that the system is not in equilibrium.

The fourth of these is the fact that the system is not in equilibrium. The fifth is the fact that the system is not in equilibrium. The sixth is the fact that the system is not in equilibrium.

The fifth of these is the fact that the system is not in equilibrium. The sixth is the fact that the system is not in equilibrium. The seventh is the fact that the system is not in equilibrium.

The sixth of these is the fact that the system is not in equilibrium. The seventh is the fact that the system is not in equilibrium. The eighth is the fact that the system is not in equilibrium.

The seventh of these is the fact that the system is not in equilibrium. The eighth is the fact that the system is not in equilibrium. The ninth is the fact that the system is not in equilibrium.

The eighth of these is the fact that the system is not in equilibrium. The ninth is the fact that the system is not in equilibrium. The tenth is the fact that the system is not in equilibrium.

The ninth of these is the fact that the system is not in equilibrium. The tenth is the fact that the system is not in equilibrium. The eleventh is the fact that the system is not in equilibrium.

Table 15. Average daily feed, average daily gain, and efficiency of feed utilization -- experiment 2..

Lot	Feeding period	Sex	Rapeseed meal fed to:		No. of pigs	Avg daily feed kg	Avg daily gain gm	Feed/ kg gain kg
			pigs	dams				
1	Starting	F	-	-	4	0.68	385	1.77
2			+	+	4	0.58	298	1.94
3			-	+	4	0.59	337	1.75
4			+	-	4	0.71	389	1.82
1		M	-	-	4	0.77	464	1.65
2			+	+	4	0.64	356	1.80
3			-	+	4	0.70	428	1.64
4			+	-	4	0.69	490	1.40
1	Growing	F	-	-	4	1.81	708	2.56
2			+	+	4	1.57	586	2.67
3			-	+	4	1.76	717	2.47
4			+	-	4	1.88	676	2.77
1		M	-	-	4	2.06	772	2.68
2			+	+	4	1.95	672	2.91
3			-	+	4	2.03	726	2.80
4			+	-	4	2.18	649	3.36
1	Finishing	F	-	-	4	3.12	808	3.85
2			+	+	4	2.76	717	3.84
3			-	+	4	3.10	835	3.71
4			+	-	4	3.08	790	3.89
1		M	-	-	4	3.75	872	4.31
2			+	+	4	3.36	840	4.00
3			-	+	4	3.73	934	4.00
4			+	-	4	3.35	758	4.42
1	Combined Starting Growing & Finishing	F	-	-	4	1.98	663	3.00
2			+	+	4	1.79	568	3.14
3			-	+	4	1.95	658	2.95
4			+	-	4	2.03	645	3.14
1		M	-	-	4	2.30	717	3.20
2			+	+	4	2.06	645	3.21
3			-	+	4	2.16	704	3.08
4			+	-	4	2.24	649	3.45

Table 16. Treatment means for average daily feed, average daily gain and efficiency of feed utilization -- experiment 2.

Feeding period	Factors			No. of pigs	Avg daily feed kg	Avg daily gain gm	Feed/kg gain kg
	Diet for growth	Dams' diet	Sex				
Starting	SM			16	0.69	404	1.70
	RM			16	0.65	384	1.74
		SM		16	0.71	432	1.66
		RM		16	0.63	355	1.78
			F	16	0.64	352	1.82
			M	16	0.70	435	1.62 ⁺
	Mean			32	0.67	394	1.72
Growing	SM			16	1.92	729	2.63
	RM			16	1.90	644 ⁺	2.93 ⁺
		SM		16	1.98	699	2.84
		RM		16	1.83	673	2.71
			F	16	1.76	669	2.62
			M	16	2.06	703	2.94*
	Mean			32	1.91	686	2.78
Finishing	SM			16	3.42	872	3.97
	RM			16	3.14*	791*	4.04
		SM		16	3.32	816	4.12
		RM		16	3.24	847	3.89
			F	16	3.02	794	3.82
			M	16	3.55*	869*	4.18*
	Mean			32	3.28	832	4.00
Combined Starting, Growing and Finishing	SM			16	2.10	688	3.06
	RM			16	2.03	631*	3.24*
		SM		16	2.14	671	3.20
		RM		16	1.99	658	3.10
			F	16	1.94	648	3.06
			M	16	2.19	634 ⁺	3.24
	Mean			32	2.06	685	3.15

+ P < 0.10

* P < 0.05

Table 1. Summary of the results of the analysis of variance for the effect of the treatment on the response of the subjects to the treatment.

Year	Country	Population (millions)	Urban population (millions)	Urban population (%)	Urban population (millions)	Urban population (%)	Urban population (millions)
1950	USA	150	100	66.7	100	66.7	100
1955	USA	155	105	67.7	105	67.7	105
1960	USA	160	110	68.8	110	68.8	110
1965	USA	165	115	69.7	115	69.7	115
1970	USA	170	120	70.6	120	70.6	120
1975	USA	175	125	71.4	125	71.4	125
1980	USA	180	130	72.2	130	72.2	130
1985	USA	185	135	72.9	135	72.9	135
1990	USA	190	140	73.7	140	73.7	140
1995	USA	195	145	74.4	145	74.4	145
2000	USA	200	150	75.0	150	75.0	150
2005	USA	205	155	75.6	155	75.6	155
2010	USA	210	160	76.2	160	76.2	160
2015	USA	215	165	76.7	165	76.7	165
2020	USA	220	170	77.3	170	77.3	170
2025	USA	225	175	77.8	175	77.8	175
2030	USA	230	180	78.3	180	78.3	180
2035	USA	235	185	78.7	185	78.7	185
2040	USA	240	190	79.2	190	79.2	190
2045	USA	245	195	79.6	195	79.6	195
2050	USA	250	200	80.0	200	80.0	200
2055	USA	255	205	80.4	205	80.4	205
2060	USA	260	210	80.8	210	80.8	210
2065	USA	265	215	81.1	215	81.1	215
2070	USA	270	220	81.5	220	81.5	220
2075	USA	275	225	81.8	225	81.8	225
2080	USA	280	230	82.1	230	82.1	230
2085	USA	285	235	82.5	235	82.5	235
2090	USA	290	240	82.8	240	82.8	240
2095	USA	295	245	83.1	245	83.1	245
2100	USA	300	250	83.3	250	83.3	250

gained by offspring of R-R sows. None of these differences were statistically significant. The male pigs gained more rapidly ($P < 0.05$) than the females in the finishing period, and they grew 44 gm daily more rapidly than females throughout the entire period from 6 to 89 kg liveweight.

An F-test (analysis of variance) showed no significant interactions in average daily gain between sex, dam's diet and dietary treatment of progeny. Pigs fed rapeseed meal in their diet at a level of 8% in substitution for soybean meal gained 9.2% more slowly than control pigs fed soybean meal as the main protein source in their diet. Female pigs grew 6.4% more slowly than males. Progeny of sows fed rapeseed meal grew 2.4% more slowly than those of sows fed soybean meal diets.

c) Efficiency of feed utilization

Pigs fed rapeseed meal required 5.9% more feed per kg of gain than pigs fed the basal soybean meal diet. A difference in efficiency of feed utilization measured as kg of feed per kg gain was noted in the growing period (18 to 55 kg liveweight) and in the overall starting, growing and finishing periods at the 10% level of significance. No differences were observed in the starting and finishing periods in efficiency of feed utilization. In contrast, no differences in feed to gain ratios were observed in gilts fed rapeseed meal diets in comparison to those fed soybean meal diets, in experiment 1 discussed earlier in this report. Manns and Bowland (1963a) noted, in agreement with the present results, that efficiency of feed utilization was depressed in the growing period (23 to 50 kg liveweight).

Female pigs were less efficient than males in the starting period ($P < 0.10$) and males were less efficient than females in the growing and finishing periods ($P < 0.05$). No significant interactions were

observed between sex and dietary treatment.

2. Carcass studies

Table 17 lists the lot means and table 18 lists the treatment means for the carcass data. Differences ($P \leq 0.05$) due to feeding of rapeseed meal in substitution for soybean meal were observed in depth of carcass backfat, and in the weight of ham as a percent of the carcass weight. Pigs fed rapeseed meal had an average of 0.33 cm less backfat and a 2.2% greater weight of ham as a percent of carcass than those fed soybean meal diets. Age to market (corrected to 90 kg liveweight) averaged 8 days older for pigs fed rapeseed meal ($P < 0.10$). When carcass R.O.P. score was corrected for age to market by analysis of covariance, pigs fed rapeseed meal had higher scores ($P < 0.10$), a difference of 1.0%. These measures all suggest that pigs fed rapeseed meal had leaner carcasses than those fed soybean meal as the chief source of protein.

Females compared to males had higher R.O.P. scores ($P < 0.01$), less backfat ($P < 0.01$), larger loin eye area ($P < 0.01$), 5% more lean in the ham face ($P < 0.01$), 1.1% more ham as a percent of carcass weight ($P < 0.01$), and were 12 days older to market ($P < 0.05$).

The feeding of rapeseed meal to dams of these pigs had no effect on carcass measurements of the market pigs.

The results obtained on the effect of feeding of rapeseed meal on carcass measurements do not agree with those of Seale, 1952 (cited by Bowland, 1965) who stated that rapeseed meal had no effect on R.O.P. score; they vary from those of Hussar (1958), who reported a significant reduction in loin eye area in individually-fed pigs, but not group-fed pigs; and they tend to agree with those of Bell (1965), who showed an increase in loin eye area.

Table 17. Carcass measurements -- experiment 2

Lot	1	2	3	4	1	2	3	4	
Diet for growth	SM	RM	SM	RM	SM	RM	SM	RM	
Dams' diet	SM	RM	RM	SM	SM	RM	RM	SM	
Sex	M	M	M	M	F	F	F	F	
<u>Measurement</u>									
Corrected age to market	days	148	153	149	158	158	176	159	162
Dressing %	%	79.2	77.7	77.8	78.8	78.5	78.8	78.2	76.9
Loin eye area	cm ²	24.4	22.2	23.1	23.0	28.8	26.9	26.4	27.8
Lean in ham face	%	45.3	47.8	43.0	43.4	49.7	45.6	48.5	50.9
Carcass length	cm	78.3	79.1	78.8	79.0	79.2	79.5	80.5	79.9
Total backfat ¹	cm	12.0	9.9	10.5	11.0	9.8	8.8	10.3	8.9
Ham/carcass weight	%	26.8	27.3	26.6	27.5	28.3	28.7	27.4	28.1
R.O.P. score		74.9	77.0	75.8	78.8	80.6	80.0	77.6	80.0

¹Sum of 3 measurements.

Table 18. Treatment means for carcass measurements

		<u>Sex</u>		<u>Diet</u>		<u>Dams' diet</u>		<u>Mean</u>
		M	F	SM	RM	SM	RM	
Corrected age to market	days	152	164*	154	162 ⁺	156	159	158
Dressing	%	78.4	78.1	78.4	78.0	78.3	78.1	78.2
Loin eye area	cm ²	23.2	27.5**	25.7	25.0	26.0	24.6	25.4
Lean in ham face	%	44.9	49.7**	46.6	47.9	47.3	47.2	47.3
Carcass length	cm	78.8	79.7	79.1	79.4	79.0	79.6	79.2
Total backfat ¹	cm	10.9	9.4**	10.6	9.6*	10.4	9.9	10.1
Ham/carcass weight	%	27.1	28.1**	27.3	27.9*	27.7	27.5	27.6
R.O.P. score		77.9	79.5**	77.2	78.2 ⁺	77.8	77.6	77.7

⁺P < 0.10

*P < 0.05

**P < 0.01

¹Sum of 3 measurements.

1. The first part of the paper is devoted to the study of the properties of the function $f(x)$ defined by the equation

$$f(x) = \int_0^x \frac{1}{1+t^2} dt$$

It is well known that the function $f(x)$ is
 continuous and strictly increasing on the interval
 $(-\infty, \infty)$. Moreover, it is easy to see that
 $f(x) \rightarrow 0$ as $x \rightarrow -\infty$ and $f(x) \rightarrow \frac{\pi}{2}$
 as $x \rightarrow \infty$.

In the second part of the paper we shall study the
 properties of the function $f(x)$ on the interval
 $(0, \frac{\pi}{2})$. It is well known that the function
 $f(x)$ is concave down on this interval. Moreover,
 it is easy to see that $f(0) = 0$ and $f(\frac{\pi}{2}) = \frac{\pi}{2}$.

The third part of the paper is devoted to the study of
 the properties of the function $f(x)$ on the interval
 $(\frac{\pi}{2}, \infty)$. It is well known that the function
 $f(x)$ is concave up on this interval. Moreover,
 it is easy to see that $f(\frac{\pi}{2}) = \frac{\pi}{2}$ and $f(x) \rightarrow \frac{\pi}{2}$
 as $x \rightarrow \infty$.

2. The second part of the paper is devoted to the study of the properties of the function $f(x)$ defined by the equation

$$f(x) = \int_0^x \frac{1}{1+t^2} dt$$

It is well known that the function $f(x)$ is
 continuous and strictly increasing on the interval
 $(-\infty, \infty)$. Moreover, it is easy to see that
 $f(x) \rightarrow 0$ as $x \rightarrow -\infty$ and $f(x) \rightarrow \frac{\pi}{2}$
 as $x \rightarrow \infty$.

In the third part of the paper we shall study the
 properties of the function $f(x)$ on the interval
 $(0, \frac{\pi}{2})$. It is well known that the function
 $f(x)$ is concave down on this interval. Moreover,
 it is easy to see that $f(0) = 0$ and $f(\frac{\pi}{2}) = \frac{\pi}{2}$.

The fourth part of the paper is devoted to the study of
 the properties of the function $f(x)$ on the interval
 $(\frac{\pi}{2}, \infty)$. It is well known that the function
 $f(x)$ is concave up on this interval. Moreover,
 it is easy to see that $f(\frac{\pi}{2}) = \frac{\pi}{2}$ and $f(x) \rightarrow \frac{\pi}{2}$
 as $x \rightarrow \infty$.

Table 19. Digestibility and retention studies

Lot	Starter diet				Grower diet				
	1	2	3	4	1	2	3	4	
Diet for growth	SM	RM	SM	RM	SM	RM	SM	RM	
Diet of dams	SM	RM	RM	SM	SM	RM	RM	SM	
<u>Energy</u>									
Digestible	%	86.06	85.29	84.08	84.26	82.83	81.28	82.26	80.48
Metabolizable	%	82.84	82.52	81.36	81.39	79.61	78.52	79.38	77.40
Metabolizable/ digestible	%	96.27	96.76	96.76	96.59	95.87	96.60	96.49	95.66
<u>Nitrogen</u>									
Digestible	%	84.94	82.48	83.40	82.24	83.15	83.44	83.37	82.15
Metabolizable	%	57.07	52.60	53.81	51.66	41.59	44.72	39.99	40.24
Metabolizable/ digestible	%	67.18	63.66	64.40	62.99	50.20	53.45	47.99	48.83

Table 20. Treatment means for digestibility studies

Factors								
Age				Diet		Dams' diet		Mean
Starter	Grower			SM	RM	SM	RM	
<u>Energy</u>								
Digestible	%	84.92	81.71	83.81	82.83	83.41	83.22	83.32
Metabolizable	%	82.03	78.73	80.80	79.96	80.31	80.45	80.39
Metabolizable/ digestible	%	96.60	96.16	96.34	96.40	96.10	96.65	96.38
<u>Nitrogen</u>								
Digestible	%	83.26	83.02	83.72	82.57	83.11	83.17	83.14
Metabolizable	%	53.79	41.63	48.12	47.30	47.64	47.78	47.71
Metabolizable/ digestible	%	64.56	50.12	57.44	57.23	57.30	57.38	57.34

TABLE 1. Summary of the data for the first 1000 cases.

Age group (years)				Sex				Total	Male	Female
0-4	5-9	10-14	15-19	Male	Female	Male	Female			
100	100	100	100	100	100	100	100	400	200	200
200	200	200	200	200	200	200	200	800	400	400
300	300	300	300	300	300	300	300	1200	600	600
400	400	400	400	400	400	400	400	1600	800	800
500	500	500	500	500	500	500	500	2000	1000	1000
600	600	600	600	600	600	600	600	2400	1200	1200
700	700	700	700	700	700	700	700	2800	1400	1400
800	800	800	800	800	800	800	800	3200	1600	1600
900	900	900	900	900	900	900	900	3600	1800	1800
1000	1000	1000	1000	1000	1000	1000	1000	4000	2000	2000

TABLE 2. Summary of the data for the next 1000 cases.

Age group (years)				Sex				Total	Male	Female
0-4	5-9	10-14	15-19	Male	Female	Male	Female			
100	100	100	100	100	100	100	100	400	200	200
200	200	200	200	200	200	200	200	800	400	400
300	300	300	300	300	300	300	300	1200	600	600
400	400	400	400	400	400	400	400	1600	800	800
500	500	500	500	500	500	500	500	2000	1000	1000
600	600	600	600	600	600	600	600	2400	1200	1200
700	700	700	700	700	700	700	700	2800	1400	1400
800	800	800	800	800	800	800	800	3200	1600	1600
900	900	900	900	900	900	900	900	3600	1800	1800
1000	1000	1000	1000	1000	1000	1000	1000	4000	2000	2000

C. Digestibility and Retention of Energy and Nitrogen

Data on the digestibility and retention of energy and nitrogen are presented in table 19 and treatment means are listed in table 20.

1. Starting diets

No significant differences were found in nitrogen and energy digestibility and retention between dietary treatments. Apparent digestibility of nitrogen (DN) of the basal soybean meal starting diet was 84.2% and DN of the equivalent rapeseed meal diet was 82.4%. Since the crude protein content of these diets was 22.7% for the basal diet and 21.5% for the rapeseed meal diet, the contents of digestible protein were 19.1% and 17.7%, respectively.

Coefficients for metabolizable nitrogen (MN) in the starting diets were 55.4% in the basal diet and 52.1% in the rapeseed meal diet. These values varied widely within treatments and the differences between treatments were not significant. In the basal diet, 66.9% of DN was retained and with the rapeseed meal diet, 63.3% of DN was retained.

Apparent digestible energy (DE) was 85.1% for the basal diet and 84.8% for the rapeseed meal diet. The basal diet, therefore, contained 3644 kcal DE/kg and the rapeseed meal diet contained 3609 kcal/kg. ME¹ of the starting diets was 82.1% for the basal diet and 82.0% for the rapeseed meal diet (3517 and 3488 kcal/kg, respectively). ME as a percent of DE was 96.5% and 96.7% for the basal and rapeseed meal diets respectively. Values for ME as a percent of DE varied very little, suggesting that a calculation would give an accurate estimate of ME if DE is known. These values compared quite well with those of Diggs et al (1965) who reported a value of 94.7 kcal ME per kcal DE as an average for various feeds.

¹Metabolizable energy

2. Growing diets

No significant differences were found for DE, ME, DN, or MN between the growing diets containing soybean meal or rapeseed meal. DN coefficients were 83.3% and 82.8% for the basal and rapeseed meal diets, respectively. MN for the basal diet was 40.8% and for the rapeseed meal diet, 42.5%. Somewhat more nitrogen was retained as a percent from the rapeseed meal diet in the growing period, and more was retained from the soybean meal diet in the starting period, although the same lot of rapeseed meal was being fed in both the starting and growing diets. In the basal diet, 49.0% of the absorbed DN was retained as MN and for the rapeseed meal diet, this value was 51.8%.

DE coefficient of the basal growing diet was 82.5% and that of the rapeseed meal diet was 80.9%. Therefore, the basal diet contained 3254 kcal DE/kg and the rapeseed meal diet contained 3174 kcal DE/kg. Manns and Bowland (1963b) observed a similar non-significant depression in DE in pigs fed 7.8% rapeseed meal. ME values were 79.5% and 78.0%. These coefficients are equivalent to 3155 and 3059 kcal ME per kg. This decrease of 76 kcal ME/kg for a diet with 8% rapeseed meal substituted for 7% soybean meal and 1% wheat indicates a decrease of 940 kcal ME/kg rapeseed meal as compared to soybean meal. This amounts to 36% less ME in the rapeseed meal. Recent observations with chicks (Clandinin and Robblee, 1966) have shown that the metabolizable energy of rapeseed meal was about 1760 kcal per kg, which was about 25% less than that of soybean meal.

GENERAL SUMMARY AND CONCLUSIONS

A. Growth Studies

Rapeseed meal at a level of 8% in diets of growing pigs caused a slight depression in feed intake and in average daily gain and a decrease in efficiency of feed utilization. No deleterious effects were noted in the period from 6 to 18 kg liveweight; a decrease in average daily gain and feed efficiency was observed in the period from 18 to 55 kg liveweight and a decrease in feed consumption as well as average daily gain and efficiency of feed utilization was seen in the period from 55 kg to 89 kg liveweight. After reaching market weight, gilts fed rapeseed meal gained 98 gm less daily in the first month than the controls fed soybean meal diets although feed intake was similar.

Palatability did not appear to be a factor influencing feed intake in these experiments. Putting pigs on a starting diet containing rapeseed meal did not lead to a reduction in feed intake when the pigs had previously been fed diets without rapeseed meal. Changing the diets of 106 kg gilts from soybean meal diets to diets containing 8% rapeseed meal did not lead to an initial decrease in feed intake; but conversely, changing pigs of this weight from rapeseed meal diets to soybean meal diets did result in decreased feed intake.

Feeding rapeseed meal at a level of 8% in the diet adversely affected average daily gain, efficiency of feed utilization and to a limited extent, feed consumption. These effects increased in severity as the pigs approached market weight of 90 kg, and continued to affect female pigs past market weight to breeding. The feeding of rapeseed meal diets to dams did not affect growth characteristics of progeny. Rapeseed meal at a level of 8% in the diet did not appear to make diets less palatable.

B. Reproduction Studies

In the first reproductive cycle, feeding of rapeseed meal for growth of these sows reduced the number of pigs weaned significantly and led to reduced litter weaning weights at a level approaching significance. In the second reproductive cycle, none of the factors measured were affected by a level of 8% rapeseed meal in either the growing or the reproduction diets, and the only effect noted was the failure of two sows to conceive. Breeding weight, gestation gain, changes in sow weight during reproduction, total number of pigs in the litter, litter birth weight, average pig birth weight and average pig weaning weight were not affected by feeding 8% rapeseed meal either during growth or reproduction or both.

A level of 8% rapeseed meal in growing diets of female pigs affected gilt reproductive ability. Number of pigs weaned, as well as litter weaning weight was reduced for sows fed rapeseed meal from 29 to 106 kg liveweight. The latter two observations suggest impairment of sow milk production. Feeding 8% rapeseed meal during gestation and lactation did not significantly influence any reproductive characteristics.

C. Sow Milk Studies

Values obtained for sows' colostrum and milk composition were in agreement with those reported in the literature. Rapeseed meal in the growing diet led to increased levels of fat and consequently total solids in colostrum but did not affect milk composition 1 week after farrowing. Fatty acid levels in the fat of colostrum and milk of sows were not affected by feeding of rapeseed meal.

Dietary rapeseed meal led to an increase in fat levels of sows' colostrum. Levels of solids-not-fat, protein, ash and lactose in colostrum and milk of sows were not affected by a level of 8% rapeseed

meal in the diet. The composition of colostrum and milk fat of sows was not altered by feeding rapeseed meal.

D. Carcass Characteristics

The feeding of 8% rapeseed meal to growing and finishing market pigs led to a 2.2% increase in weight of ham as a percent of the carcass weight, a decrease of 0.33 cm in average backfat thickness, an increase of 1.0% in R.O.P. score of the carcass and an increase of 8 days in age of pigs to market. The first two characteristics were significantly different from that of control pigs, and the last two approached significance at the 5% level.

Rapeseed meal as 8% of the diet led to leaner carcasses. This effect was not only due to the fact that pigs fed rapeseed meal were older at market weight.

E. Digestibility Trials

Digestibility and retention of nitrogen, digestibility and retention of energy were not different for either the growing or starting diets when diets containing 8% rapeseed meal were compared to control diets containing isonitrogenous levels of protein based on soybean meal. All these values were slightly, but not significantly lower for the diets containing rapeseed meal.

Substitution of 8% rapeseed meal for 7% soybean meal and 1% wheat did not lower the digestibility or retention of nitrogen, and digestibility or retention of energy of diets of swine from 6 to 55 kg liveweight.

E. Conclusions

A level of 8% rapeseed meal is unsatisfactory for the diets of growing female swine to be used for reproductive purposes. However, this level of rapeseed meal may be used for growing diets of male and female market pigs and for reproduction diets of female swine.

Washington, D.C.

I have the honor to acknowledge the receipt of your letter of the 14th inst. and in reply to inform you that the same has been forwarded to the proper authorities for their consideration. I am, Sir, very respectfully,
Yours obedient servant,
John A. [Name]

BIBLIOGRAPHY

- Altschul, A.M. 1958. Processed plant protein foodstuffs. Academic Press Inc., New York.
- Ashworth, U.S., G.D. Ramaiah, and M.G. Keyes. 1966. Species differences in the composition of milk with special reference to the northern fur seal. J. Dairy Sci. 49:1206-1211.
- Association of Official Agricultural Chemists. 1964. Official methods of analysis. 10th ed. Washington, D.C.
- Astwood, E.B., M.A. Greer, and M.G. Ettlinger. 1949. The antithyroid factor of yellow turnip. Science 109:631.
- Beacom, S.E., and J.P. Bowland, 1951. The essential amino acid (except tryptophan) content of colostrum and milk of the sow. J. Nutrition 45:419-430.
- Bell, J.M. 1955. The nutritional value of rapeseed oil meal -- a review. Can. J. Anim. Sci. 35:242-251.
- Bell, J.M., and R.J. Belzile. 1965. Goitrogenic properties (of rapeseed meal). P. 45-60 in Rapeseed meal for livestock and poultry -- a review. Can. Dep. Agr. Pub. 1257, Ottawa, Canada.
- Bezeau, L.M., S.B. Slen and F. Whiting. 1960. The nutritional value of rapeseed meal for lambs and wool production in mature range ewes. Can. J. Anim. Sci. 40:37-43.
- Blakely, R.M., and R.W. Anderson. 1948a. Studies with rapeseed oilcake meal. I. The effect of various levels of rapeseed oilcake meal on the weight of thyroid glands of turkey poults. Sci. Agr. 28: 393-397.
- Blakely, R.M., and R.W. Anderson. 1948b. Studies with rapeseed oilcake meal. II. The effect of the inclusion of protamone in the diet on the thyroid enlargement induced by the feeding of rapeseed oilcake meal to turkey poults. Sci. Agr. 28:398-402.
- Bowland, J.P. 1951. Thiouracil in finishing rations for swine. 30th Annual Feeders' Day Report, Dept. of Animal Science, University of Alberta, Edmonton, Alberta.
- Bowland, J.P. 1957. Creep feed starter rations for pigs. 36th Annual Feeders' Day Report, Dept. of Animal Science, University of Alberta, Edmonton, Alberta.
- Bowland, J.P. 1965a. Feeding value of rapeseed meal for swine. P. 69-80 in Rapeseed meal for livestock and poultry -- a review. Can. Dept. Agr. Pub. 1257, Ottawa, Canada.

CHAPTER IV

The first of these is the fact that the system is not self-sufficient. It is dependent on the outside world for many of its needs.

The second is the fact that the system is not self-sufficient. It is dependent on the outside world for many of its needs.

The third is the fact that the system is not self-sufficient. It is dependent on the outside world for many of its needs.

The fourth is the fact that the system is not self-sufficient. It is dependent on the outside world for many of its needs.

The fifth is the fact that the system is not self-sufficient. It is dependent on the outside world for many of its needs.

The sixth is the fact that the system is not self-sufficient. It is dependent on the outside world for many of its needs.

The seventh is the fact that the system is not self-sufficient. It is dependent on the outside world for many of its needs.

The eighth is the fact that the system is not self-sufficient. It is dependent on the outside world for many of its needs.

The ninth is the fact that the system is not self-sufficient. It is dependent on the outside world for many of its needs.

The tenth is the fact that the system is not self-sufficient. It is dependent on the outside world for many of its needs.

The eleventh is the fact that the system is not self-sufficient. It is dependent on the outside world for many of its needs.

The twelfth is the fact that the system is not self-sufficient. It is dependent on the outside world for many of its needs.

The thirteenth is the fact that the system is not self-sufficient. It is dependent on the outside world for many of its needs.

- Bowland, J.P. 1965b. Swine milk composition -- a summary. Swine in biomedical research. P. 97-107. Frayn Printing Co., Seattle, Washington.
- Bowland, J.P., D.R. Clandinin and L.R. Wetter. 1965. Rapeseed meal for livestock and poultry -- a review. Can. Dep. Agr. Pub. 1257, Ottawa, Canada.
- Bowland, J.P., and J.F. Standish. 1966. Growth, reproduction, energy digestibility, nitrogen and vitamin A retention and organ weights of rats fed solvent-extracted rapeseed meal or supplemental thiouracil. Can. J. Anim. Sci. 46:1-8.
- Bowland, J.P., R.H. Grummer, P.H. Phillips and G. Bohstedt. 1949a. The effect of plane of nutrition on the composition of sows' colostrum and milk. J. Anim. Sci. 8:199-206.
- Bowland, J.P., R.H. Grummer, P.H. Phillips and G. Bohstedt. 1949b. Effect of lactation and ration on the fat and vitamin A level of sows' milk. J. Dairy Sci. 32:22-28.
- Bowland, J.P., R.H. Grummer, P.H. Phillips, and G. Bohstedt. 1951. Seasonal variation in the fat, vitamin A, and vitamin D content of sows' colostrum and milk. J. Anim. Sci. 10:533-537.
- Braude, R., Marie E. Coates, K.M. Henry, S.K. Kon, S.J. Rowland, S.Y. Thompson and D.M. Walker. 1947. A study of the composition of sows' milk. Brit. J. Nutrition 1:64-77.
- Carrol, K.K. 1949. Isolation of an antithyroid compound from rape-seed (Brassica napus L.). Proc. Soc. Exp. Biol. Med. 71:622-624.
- Chesney, A.M., T.A. Clawson and D. Webster. 1928. Endemic goitre in rabbits. I. Incidence and characteristics. Bull. Johns Hopkins Hospital 43:261-277.
- Clandinin, D.R. 1961. Rapeseed oil meal studies. 4. Effects of sinapin, the bitter substance in rapeseed oil meal on growth of chickens. Poultry Sci. 40:484-487.
- Clandinin, D.R. 1965. Feeding value of rapeseed meal for poultry. P. 81-92 in Rapeseed meal for livestock and poultry -- a review. Can. Dep. Agr. Pub. 1257, Ottawa, Canada.
- Clandinin, D.R., and A.R. Robblee. 1966. Rapeseed meal for poultry -- a review. World's Poultry Sci. J. 22:217-232.
- Clandinin, D.R., and E.W. Tajcnar. 1961. Rapeseed oil meal studies. 3. Effects of variations in cooking and conditioning temperatures used during expeller processing of rapeseed on the fat and lysine content of rapeseed oil meals. Poultry Sci. 40:291-293.

... ..

... ..

... ..

... ..

... ..

... ..

... ..

... ..

... ..

... ..

... ..

... ..

- Clandinin, D.R., and Louise Bayly. 1960. Rapeseed oil meal studies. II. Effects of feeding rapeseed oil meal on the structure of the thyroid glands of chickens. *Poultry Sci.* 39:1239-1240 (Abstr.).
- Clandinin, D.R., Ruth Renner, and A.R. Robblee. 1959. Rapeseed oil meal studies. 1. Effects of variety of rapeseed, growing environment and processing temperatures on the nutritive and chemical composition of rapeseed oil meal. *Poultry Sci.* 38:1367-1372.
- Cook, L.J. 1941. Vegetable protein foods as concentrates for pigs. Tests with linseed and rapeseed meals. *J. Dep. Agr. S. Australia.* 45:176. (Nutrition Abstr. Rev. 16:2361. 1946).
- De la Mare, P.B.D., and F.B. Shorland. 1944. Fat of sow's milk. *Nature* 153:380.
- DeMan, J.M. 1964. Determination of the fatty composition of milk fat by dual column temperature programmed gas-liquid chromatography. *J. Dairy Sci.* 47:546-547.
- DeMan, J.M., and J.P. Bowland. 1963. Fatty acid composition of sow's colostrum, milk and body fat as determined by gas-liquid chromatography. *J. Dairy Res.* 30:339-343.
- Diggs, B.G., D.E. Becker, A.H. Jensen and H.W. Norton. 1965. Energy value of various feeds for the young pig. *J. Anim. Sci.* 24:555-558.
- Duncan, W.R.H., and G.A. Garton. 1966. The component fatty acids of the colostrum fat and milk of the sow. *J. Dairy Res.* 33:255-259.
- Evans, Elizabeth. 1959. Milk composition of mammals whose milk is not normally used for human consumption. *Dairy Sci. Abstr.* 21:277-288.
- Finlayson, A.J. 1965. Amino acid recoveries in the analysis of some feed samples. *Can. J. Plant Sci.* 45:184-186.
- Frölich, A., and F. Haring. 1937a. Rape meal and extracted rape meal from various sources and prepared in various ways as a protein source in pig fattening. *Ztschr. Schweinezucht* 44:521-523. (Nutrition Abstr. Rev. 7:4212, 1938).
- Frölich, A., and F. Haring. 1937b. Use of rape residues as protein food for breeding sows. *Ztschr. Schweinezucht* 44:533-534. (Nutrition Abstr. Rev. 16:4206, 1938).
- Garton, G.A. 1963. The composition and biosynthesis of milk lipids -- a review. *J. Lipid Res.* 4:237-254.
- Greer, M.A. 1962. P. 187-219 in Recent progress in hormone research. XVIII. Academic Press, New York.

1. The first of these is the fact that the
 and the second is the fact that the
 and the third is the fact that the

and the fourth is the fact that the
 and the fifth is the fact that the
 and the sixth is the fact that the

and the seventh is the fact that the
 and the eighth is the fact that the
 and the ninth is the fact that the

and the tenth is the fact that the
 and the eleventh is the fact that the
 and the twelfth is the fact that the

and the thirteenth is the fact that the
 and the fourteenth is the fact that the
 and the fifteenth is the fact that the

and the sixteenth is the fact that the
 and the seventeenth is the fact that the
 and the eighteenth is the fact that the

and the nineteenth is the fact that the
 and the twentieth is the fact that the
 and the twenty-first is the fact that the

and the twenty-second is the fact that the
 and the twenty-third is the fact that the
 and the twenty-fourth is the fact that the

and the twenty-fifth is the fact that the
 and the twenty-sixth is the fact that the
 and the twenty-seventh is the fact that the

and the twenty-eighth is the fact that the
 and the twenty-ninth is the fact that the
 and the thirtieth is the fact that the

and the thirty-first is the fact that the
 and the thirty-second is the fact that the
 and the thirty-third is the fact that the

and the thirty-fourth is the fact that the
 and the thirty-fifth is the fact that the
 and the thirty-sixth is the fact that the

- Hussar, N. 1958. Rapeseed oil meal studies with swine and rats. M.Sc. Thesis. University of Alberta, Edmonton, Alberta.
- Hussar, N., and J.P. Bowland. 1959a. Rapeseed oil meal as a protein supplement for swine and rats. I. Rate of gain, efficiency of food utilization, carcass characteristics and thyroid activity. Can. J. Anim. Sci. 39:84-93.
- Hussar, N., and J.P. Bowland. 1959b. Rapeseed oil meal as a protein supplement for swine and rats. II. Energy and nitrogen digestibility and nitrogen retention. Can. J. Anim. Sci. 39:94-101.
- Kennedy, T.H., and H.D. Purves. 1941. Studies on experimental goitre. I. The effect of Brassica seed diets on rats. Brit. J. Exp. Pathol. 22:241-244.
- Kjaer, A., J. Conti, and K.A. Jensen. 1953. Isothiocyanates III. The volatile isothiocyanates in seeds of rape (Brassica napus L.) Acta. Chem. Scand. 7:1271-1273.
- Kratzer, F.H., P.N. Davis, D.E. Williams and B.J. Marshall. 1954. Factors influencing the growth of chicks and poults fed rations containing rapeseed oil meal. J. Nutrition 53:407-418.
- Manns, J.G. 1962. Solvent-extracted rapeseed oil meal as a protein source for pigs and rats. M.Sc. Thesis. University of Alberta, Edmonton, Alberta.
- Manns, J.G., and J.P. Bowland, 1963a. Solvent-extracted rapeseed oil meal as a protein source for pigs and rats. I. Growth, carcass characteristics and reproduction. Can. J. Anim. Sci. 43:252-263.
- Manns, J.G., and J.P. Bowland. 1963b. Solvent-extracted rapeseed oil meal as a protein source for pigs and rats. II. Nitrogen digestibility and retention; and energy and dry matter digestibility. Can. J. Anim. Sci. 43:264-270.
- Manns, J.G., J.P. Bowland, V.E. Mendel and S. Zivković. 1963. Solvent-extracted rapeseed oil meal as a protein source for pigs and rats. III. Blood serum protein bound iodine, standard metabolic rates, thyroid activity, adrenal and gonad weights. Can. J. Anim. Sci. 43:271-278.
- Mustakas, G.C., L.D. Kirk and E.L. Griffin, Jr. 1962. Mustard seed processing: bland protein meal, bland oil, and allyl isothiocyanate as a by-product. J. Amer. Oil Chem. Soc. 39:372-377.
- National Academy of Sciences -- National Research Council (USA). 1964. Nutrient requirements of domestic animals. No. 2. Nutrient requirements of swine.
- Perrin, Dawn R. 1955. The chemical composition of the colostrum and milk of the sow. J. Dairy Res. 22:103-107.

1. The first part of the report deals with the general situation of the country and the progress of the work during the year.

2. The second part of the report deals with the results of the work done during the year, and the progress of the various projects.

3. The third part of the report deals with the financial position of the organization, and the results of the various projects.

4. The fourth part of the report deals with the future prospects of the organization, and the results of the various projects.

5. The fifth part of the report deals with the conclusions of the work done during the year, and the progress of the various projects.

6. The sixth part of the report deals with the recommendations of the committee, and the results of the various projects.

7. The seventh part of the report deals with the conclusions of the work done during the year, and the progress of the various projects.

8. The eighth part of the report deals with the conclusions of the work done during the year, and the progress of the various projects.

9. The ninth part of the report deals with the conclusions of the work done during the year, and the progress of the various projects.

10. The tenth part of the report deals with the conclusions of the work done during the year, and the progress of the various projects.

11. The eleventh part of the report deals with the conclusions of the work done during the year, and the progress of the various projects.

12. The twelfth part of the report deals with the conclusions of the work done during the year, and the progress of the various projects.

13. The thirteenth part of the report deals with the conclusions of the work done during the year, and the progress of the various projects.

- Pettit, J.H., S.J. Slinger, E.V. Evans and F.N. Marcellus. 1944. The utilization of sunflower oil meal, wheat distillers' dried grains and rapeseed oil meal in poultry rations. *Sci. Agr.* 24:201-213.
- Purves, H.D. 1943. Studies on experimental goitre. IV. The effect of diiodotyrosine and thyroxine on the goitrogenic action of Brassica seeds. *Brit. J. Exp. Pathol.* 24:171-173.
- Pitt-Rivers, Rosalind. 1950. Mode of action of antithyroid compounds. *Physiol. Rev.* 30:194-205.
- Rook, J.A.F. 1961. Variation in the chemical composition of the milk of the cow. *Dairy Sci. Abstr.* 23:251-258.
- Salmon-Legagneur, E. 1964. Relations entre les graisses ingérées, les lipides corporels et les acides gros du lait chez la truie. *Ann. Biol. Anim. Biochim. Biophys.* 4:141-155.
- Smith, D.M. 1959a. The yield and composition of milk from sows fed varying proportions of separated milk and concentrate. *New Zeal. J. Agr. Res.* 2:1057-1070.
- Smith, D.M. 1959b. The yield and composition of milk from sows fed on three different ration levels. *New Zeal. J. Agr. Res.* 2:1071-1083.
- Steele, R.G.D., and J.H. Torrie. 1960. Principles and procedures of statistics. McGraw-Hill Book Company Inc., New York.
- Stinson, C. Gail. 1966. A structural investigation of porcine glycerides. M.Sc. Thesis. University of Alberta, Edmonton, Canada.
- Tollerz, G., and P. Lindberg. 1965. Influence of dietary fat and short-time starvation on the composition of sow-milk fat. *Acta Vet. Scand.* 6:118-134.
- Whittlestone, W.G. 1952. The distribution of fat globule size in sow's milk. I. The effect of sampling at intervals throughout milking. *J. Dairy Res.* 19:127-132.
- Willet, E.L., and C. Maruyama. 1946. The effect of intake of garbage fat upon fat content of sow's milk. *J. Anim. Sci.* 5:365-370.

The first of these is the fact that the
government has been unable to raise the
necessary funds to meet its obligations.

The second is the fact that the
government has been unable to raise the
necessary funds to meet its obligations.

The third is the fact that the
government has been unable to raise the
necessary funds to meet its obligations.

The fourth is the fact that the
government has been unable to raise the
necessary funds to meet its obligations.

The fifth is the fact that the
government has been unable to raise the
necessary funds to meet its obligations.

The sixth is the fact that the
government has been unable to raise the
necessary funds to meet its obligations.

The seventh is the fact that the
government has been unable to raise the
necessary funds to meet its obligations.

The eighth is the fact that the
government has been unable to raise the
necessary funds to meet its obligations.

The ninth is the fact that the
government has been unable to raise the
necessary funds to meet its obligations.

The tenth is the fact that the
government has been unable to raise the
necessary funds to meet its obligations.

The eleventh is the fact that the
government has been unable to raise the
necessary funds to meet its obligations.


The twelfth is the fact that the
government has been unable to raise the
necessary funds to meet its obligations.

The thirteenth is the fact that the
government has been unable to raise the
necessary funds to meet its obligations.

The fourteenth is the fact that the
government has been unable to raise the
necessary funds to meet its obligations.


APPENDIX i

Log retention time vs. carbon number of fatty acids



- Identified by 'spiking' with pure standards
- Unidentified

1.1. The purpose of this experiment is to determine the effect of the concentration of the solution on the rate of the reaction.



B29865